

## Correspondence

# Primary fallopian tube carcinoma

To the Editors:

We read the report by Ou et al entitled “Primary fallopian tube carcinoma: clinicopathological analysis of 12 cases” with interest [1]. At least 22 studies have addressed this topic, as reviewed by the authors [1]. The authors provided a reasonable conclusion that a diagnosis of primary fallopian tube carcinoma (PFTC) should be considered in patients who complain of lower abdominal pain in association with vaginal bleeding and/or watery discharge (16.7%) or tubo-ovarian abscess (TOA) (25%) [1].

Although this conclusion from the authors might be correct, we do not think that this concept could be successfully applied in our routine practice. Of course, we would like to emphasize that this should not be construed as an argument against the authors’ excellent work.

First, the incidence of TOA might not be very high, but it is definitely far higher than that of PFTC [2,3]. Therefore, it is potentially dangerous and impractical to conclude that the diagnosis of TOA should consider the possibility of PFTC. In addition, pelvic inflammatory disease and its severe form—TOA—are medical illnesses, and the usual choice of treatment is conservative medical treatment with antibiotics, not surgery [4,5]. In addition, if surgery is recommended, the laparoscopic approach might be a good alternative to traditional surgery for managing patients with TOA because it can be easily performed by a qualified gynecologist and is an acceptable treatment for most patients [6]. Furthermore, it is sometimes difficult to make a differential diagnosis of TOA if other uncertain conditions also present (such as corpus luteum rupture with hemorrhage) and the patient only complains of dull pain [7]. In addition, and most importantly, the laparoscopic approach is often considered for treating these uncertain conditions before performing more the invasive conventional laparotomy [8,9]. We wonder if the authors performed any laparoscopic surgeries on these “supposed” TOA patients or not. In addition, we encourage the authors to report the age distribution of these three patients with “supposed” TOA and if these patients were treated with antibiotics.

In summary, it is difficult to know whether or not the presence of abdominal pain in a patient with PFTC is highly significant, as with the diagnosis of ovarian neoplasms or other uncertain abdominal conditions. The two conditions are

always controversial, although keeping in mind the possibility of any life-threatening or severe diseases, as suggested by Ou et al, is always encouraged [10].

## References

- [1] Ou YC, Huang HY, Huang CC, Chang Chien CC, Tseng CW, Lin H. Primary fallopian tube carcinoma: clinicopathological analysis of 12 cases. *Taiwan J Obstet Gynecol* 2011;50:141–4.
- [2] Wang PH, Yuan CC, Chao HT, Juang CM, Ng HT. Prognosis of primary fallopian tube adenocarcinoma: report of 25 patients. *Eur J Gynaecol Oncol* 1998;19:571–4.
- [3] Wang PH, Yuan CC, Chao HT. Surgico-pathological stage I primary fallopian tube adenocarcinoma. *Int J Gynaecol Obstet* 1998;63:73–4.
- [4] Su WH, Tsou TS, Chen CS, Ho TY, Lee WL, Yu YY, et al. Are we satisfied with the tools for the diagnosis of gonococcal infection in females? *J Chin Med Assoc* 2011;74:430–4.
- [5] Su WH, Tsou TS, Chen CS, Ho TY, Lee WL, Yu YY, et al. Diagnosis of Chlamydia infection in females. *Taiwan J Obstet Gynecol* 2011;50:261–7.
- [6] Yang CC, Chen P, Tseng JY, Wang PH. Advantages of open laparoscopic surgery over exploratory laparotomy in patients with tubo-ovarian abscess. *J Am Assoc Gynecol Laparosc* 2002;9:327–32.
- [7] Wang PH, Yen MS. Primary ovarian pregnancy. *Taiwan J Obstet Gynecol* 2008;47:377–8.
- [8] Huang BS, Wang PH. Ovarian torsion during pregnancy. *Taiwan J Obstet Gynecol* 2011;50:409–10.
- [9] Ding DC, Chu TY. Disseminated granulosa cell tumor of pelvis. *Taiwan J Obstet Gynecol* 2011;50:236–9.
- [10] Horng HC, Wang PH. Ovarian cancer presenting as an acute abdomen was successfully diagnosed and managed by laparoscopy. *Taiwan J Obstet Gynecol*, in this issue.

Ben-Shian Huang  
Kuan-Chong Chao

*Department of Obstetrics and Gynecology,  
National Yang-Ming University School of Medicine,  
Taipei, Taiwan  
National Yang-Ming University Hospital, Ilan, Taiwan  
Department of Obstetrics and Gynecology,  
Taipei Veterans General Hospital, Taipei, Taiwan*

Wen-Hsun Chang  
*Department of Nursing, Taipei Veterans General Hospital,  
Taipei, Taiwan  
Institute of Hospital and Health Care Administration,  
National Yang-Ming University School of Medicine,  
Taipei, Taiwan*

Peng-Hui Wang\*

*Department of Obstetrics and Gynecology,  
National Yang-Ming University School of Medicine,  
Taipei, Taiwan  
National Yang-Ming University Hospital, Ilan, Taiwan  
Department of Obstetrics and Gynecology,  
Taipei Veterans General Hospital, Taipei, Taiwan*

\*Corresponding author. Department of Obstetrics and  
Gynecology, National Yang-Ming University,  
Taipei Veterans General Hospital and  
National Yang-Ming University Hospital,  
201, Section 2, Shih-Pai Road, Taipei 112, Taiwan.  
E-mail addresses: [phwang@vghtpe.gov.tw](mailto:phwang@vghtpe.gov.tw),  
[phwang@ym.edu.tw](mailto:phwang@ym.edu.tw)