

Review Article

Curcuminoids and resveratrol as anti-Alzheimer agents

Oliver B. Villaflores^a, Ying-Ju Chen^b, Chih-Ping Chen^{e,f,g,h,i,j}, Jui-Ming Yeh^{a,**},
Tzong-Yuan Wu^{b,c,d,*}^aDepartment of Chemistry, Chung Yuan Christian University, Chung Li, Taiwan^bDepartment of Bioscience Technology, Chung Yuan Christian University, Chung Li, Taiwan^cCenter for Nanotechnology, Chung Yuan Christian University, Chung Li, Taiwan^dR & D Center for Membrane Technology, Chung Yuan Christian University, Chung Li, Taiwan^eDepartment of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan^fDepartment of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan^gDepartment of Biotechnology, Asia University, Taichung, Taiwan^hSchool of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, TaiwanⁱInstitute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan^jDepartment of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

Accepted 20 July 2012

Abstract

Alzheimer disease (AD) is by far the most common cause of dementia globally. This neurodegenerative disorder of the brain is chronic and progressive, characterized clinically by the deterioration in the key symptoms of behavioral and cognitive abilities. Treatment options for this disease currently are limited. Deposition of amyloid- β and tau hyperphosphorylation are cardinal pathologic features of AD that lead to the formation of neuronal plaques and neurofibrillary tangles, respectively. In addition to mounting research on herbal compounds for the treatment of AD, curcuminoids and resveratrol appear to be beneficial as anti-AD agents. Curcuminoids (curcumin and demethoxycurcumin) and resveratrol possess unique properties that make them especially worthy of further studies. This review article revisits and presents the current research done on the potential of the curcuminoids curcumin and demethoxycurcumin and the polyphenolic compound resveratrol as anti-AD compounds.

Copyright © 2012, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

Keywords: Alzheimer's disease; amyloid- β ; curcuminoids; resveratrol; tau hyperphosphorylation

Introduction

Demographic ageing is a global process that denotes the success of improved healthcare over the past years. Today, human life expectancy is significantly longer resulting to a much greater proportion of elderly people. Nevertheless, ageing does impose some challenges as well. It has been well

documented that corollary to ageing there is also an increase in cases of Alzheimer's disease and other dementias in the senile but modern society. It is the aim of this review paper to present the benefits and potentials of curcumin, demethoxycurcumin and resveratrol as a therapeutic agent for Alzheimer's disease.

Alzheimer disease—an emerging burden on society

The globally prevalent illness Alzheimer disease (AD) is a genetically complex disorder that accounts for most cases of dementia experienced by older people. It is a neurodegenerative disorder affecting major brain areas including the cortex and limbic system, and is characterized by progressive decline

* Corresponding author. Department of Bioscience Technology, Chung Yuan Christian University, Chung Li, Taiwan.

** Corresponding author. Department of Chemistry, Chung Yuan Christian University, Chung Li, Taiwan.

E-mail addresses: juiming@cycu.edu.tw (J.-M. Yeh), tywu@cycu.edu.tw (T.-Y. Wu).

in memory and impairment of at least one other cognitive function [1]. AD often begins with symptoms such as short-term memory loss and progresses with more widespread cognitive and emotional dysfunction. The so-called late-onset AD occurs after age 65 years. AD features ongoing deterioration of the patient's functioning, which results in substantial and long-lasting disability over the approximate 7–10 years from diagnosis to eventual death [2]. A diagnosis of AD is correct when the disorder is not limited to the period of delirium, is present for at least 6 months, and has no alternative explanation [3].

The health and social welfare of elderly people have long been underprioritized in the public health policy of most countries. The progressive mental deterioration in old age has been recognized and described throughout history. However, it was not until 1906 that a German physician, Alois Alzheimer, identified a collection of brain cell abnormalities as a disease [4]. He presented in his paper the results of postmortem studies on his patient, a 51-year-old woman named Auguste D. who had experienced a progressive presenile dementia [5], the first case of AD ever documented. In Dr. Alzheimer's paper, the patient showed strong feelings of jealousy toward her husband, followed by rapid memory impairment and disorientation in time and space. She then became bedridden and incontinent. She died 4.5 years after the onset of the illness. Her brain autopsy revealed an evenly affected atrophic brain without microscopic foci. The larger cerebral vessels showed arteriosclerotic changes [6]. Dr. Alzheimer also observed the presence of usual deposits in the cortex that were refractory to staining.

Dementia has contributed 11.2% of years lived with disability in elderly people age 60 years and older; more than stroke (9.5%), musculoskeletal disorders (8.9%), cardiovascular disease (5.0%), and all forms of cancer (2.4%) as reported in the global burden of diseases of the 2003 World Health Report. Based on the 2005 Lancet study by Ferri et al [7] on the global prevalence of dementia, 4.6 million new cases of dementia are projected to be recorded each year, with a rate of one case per 7 seconds, and the number of people affected will double every 20 years to as high as 81.1 million by 2040. It was predicted that most people with dementia will live in developing countries (60% in 2001, rising to 71% by 2040). However, the rates of increase are not uniform; thus, numbers in developed countries are forecasted to increase by 100% between 2001 and 2040, but more than 300% in India, China, and their South Asian and Western Pacific neighbors. In addition, there are currently an estimated 18 million people worldwide in whom AD has been diagnosed, according to the World Health Organization. In 2025, it is estimated that this figure will double to approximately 34 million [8]. Although people with dementia are heavy consumers of health services, direct costs in developed countries occur mostly from community and residential care. Despite technologic advancements in medicine, research revealed that most people currently living with dementia have not received a formal diagnosis. In countries with high income, only 20–50% of dementia cases are recognized and

are documented during primary care. This treatment gap is without a doubt much greater in low- and middle-income countries, with one study in India suggesting 90% remain unidentified [9]. If the current statistics are extrapolated to other countries worldwide, it suggests that approximately 28 million of the 36 million people with dementia have not received a diagnosis and therefore do not have access to the treatment, care, and organized support that a formal diagnosis can provide. This situation clearly presents a major concern, given that the world's population is aging; new cases of dementia and AD are steadily increasing. Therefore, early diagnosis and intervention are important mechanisms by which the treatment gap can be closed.

Amyloid β —the culprit in AD pathology

In past years, important progress has been made in the understanding of the pathogenic mechanisms of AD, and new therapeutic drugs have become available that allowed the underlying disease process to be treated directly. In this respect, the “amyloid hypothesis” has become the dominant theory in this field. It is believed that amyloid beta ($A\beta$) accumulation in plaques or as partial soluble filaments initiates a pathologic cascade leading to tangle formation [10], neuronal dysfunction, and possibly inflammation and oxidative damage, with neurodegeneration and dementia as the final outcome. The $A\beta$ is a 40–42 amino acid peptide that has a β -pleated sheet amyloidogenic nature, as detected by Congo red and thioflavin-S staining. The $A\beta$ peptides, which are 3–4 kDa in length, are the building blocks of both the amyloid fibrils in neuritic plaques, in addition to the vascular deposits that accumulate in the brains of AD patients. The $A\beta$ is derived from the proteolytic processing of one or more isoforms of the amyloid- β precursor protein (APP) [11]. Furthermore, the $A\beta$ can exist in two primary forms: as a soluble peptide or in an aggregated state as insoluble amyloid deposits in a beta-sheet conformation; this aggregated state forms the basis of the senile plaques characteristic of AD. Previous studies have shown the neurotrophic characteristic of $A\beta$ at low dosages (10^{-10} M to 10^{-8} M); nevertheless, at higher doses ($>10^{-7}$ M) the peptide becomes neurotoxic [12].

There are also some reports that the protofibrils of $A\beta$, which are metastable intermediates in amyloid fibril formation, may be toxic [13]. The size of $A\beta$ oligomers is distributed over a wide molecular weight range (from <10 kDa to >100 kDa), with structural polymorphism in $A\beta$ oligomers of similar sizes. The soluble $A\beta$ oligomers are found to be more cytotoxic than fibrillar $A\beta$ aggregates in general and inhibit many critical neuronal activities including long-term potentiation, a classic model for synaptic plasticity and memory loss *in vivo* and in culture [14,15]. Low-molecular-weight $A\beta$ oligomers found in the soluble fractions of the human brain and amyloid plaque extracts [16] were suggested to be the building blocks of larger oligomers or insoluble amyloid fibrils [17,18]. The high toxicity of low-molecular-weight $A\beta$ -oligomers is also supported by *in vitro* studies showing that $A\beta$ dimers are threefold more toxic than monomers and that tetramers are 13-fold more

toxic [16]. Synthetic A β s were also found to be toxic because these can form fibrillar aggregates that have properties similar to those found in AD plaques in the brain. Lambert et al [14] published the formation of small globular A β oligomers (approximately 5 nm in diameter) in Ham's-F12 medium. These globular oligomers were referred as A β -derived diffusible ligands. These molecules strongly bind with the dendritic arbors of cultured neurons to cause neuronal death and the blocking of long-term potentiations [14].

Upon reaching the critical concentration, with the A β aggregates forming insoluble amyloidogenic deposits, there is then an observed general age-related increase in A β generation by the neural cells [19]. Studies have shown that the 42-amino-acid form of the A β is the more amyloidogenic of the two forms of the peptide, and it manifests a greater tendency to create insoluble deposits commonly seen in brains affected by AD [20–22]. In certain familial AD, including cases of APP mutation [23] and presenilin mutations [24], there is an increase in the A β_{42} :A β_{40} , with the larger form of the peptide acting as a potential “seed” for the deposition of the amyloid peptide and plaque formation. The deposition of the A β appeared to be an early event in AD-associated neurodegeneration, with this peptide acting as a seed for the subsequent deposition of the A β_{40} peptide.

Amyloid- β precursor protein—a multifunctional molecule in the central nervous system

Although there are many components contained within the neuritic plaques of the brain affected by AD, the APP has a distinct role toward the pathogenesis of AD. The APP isoforms are Type I transmembrane sialoglycoproteins encoded for by a single gene on chromosome 21 that contains 19 exons [25]. The three primary APP isoforms are as follows: APP₆₉₅, APP₇₅₁, and APP₇₇₀. The APP is a multifunctional protein that serves in both cell adhesion and neurite outgrowth. The APP protein has been demonstrated to interact with a number of elements of the extracellular matrix that include collagens I and IV [26], laminin [27], fibronectin [28], heparan sulfate [29], and some glycosaminoglycans [30]. APP has also been proposed to play a role in the modulation of synaptic plasticity [31]. This protein can also play a role in long-term potentiation [32]. The soluble form of APP may also elicit both neurotrophic and neuroprotective properties. The APP acts as a neuroprotective agent, stabilizing intracellular calcium levels wherein the treatment of cells with APP resulted in rapid and prolonged decrease in cellular Ca²⁺ [33]. The APP was able to attenuate the toxic actions of the A β peptide by overcoming the A β -associated elevation of free intracellular Ca²⁺ levels with the associated induction of free radical species [34]. As a neurotrophic agent, APP stimulates neuronal cell differentiation and neurite outgrowth. Transfection of an APP-null neural cell with APP complementary DNA (cDNA) resulted in an increased neurite outgrowth. The binding of a larger peptide (Ala₃₁₉ to Met₃₃₅) to cell membrane preparation is both saturable and specific, suggesting that soluble APPs (APPs) mediates its effect by a receptor-mediated mechanism [35].

With regard to APP protein processing, two alternate pathways are known. The primary pathway produces APP after the cleavage at Lys¹⁶ within the A β region of the protein by the enzyme, alpha (α)-secretase. The alternative pathway can be through the cleavage of the APP protein by the β - and γ -secretase to generate the 40–42 amino acid A β peptide. The major species of the A β produced at normal circumstances is the 40 amino acid form (A β_{40}). However, A β_{42} was found to be increased in AD and was observed to be more amyloidogenic and acts as a seed for amyloid deposition [36]. The pathway by which APP is processed can be influenced by a multitude of factors that may be extracellular or intracellular. In addition, the processing mechanisms could be cell-, tissue-, or species-specific. There is also an increasing number of evidence that the carboxy-terminal fragment of the APP may also exhibit neurotoxicity in addition to A β [37]. The toxic fragment was identified as a 100-amino-acid C-terminal fragment (C100 protein). This fragment inhibits the Mg²⁺/Ca²⁺-adenosine triphosphatase within the endoplasmic reticulum and also the Na⁺/Ca²⁺ exchanger. This in turn compromises the ability of the cells to sequester Ca²⁺ within the endoplasmic reticulum, thereby raising the level of the intracellular Ca²⁺ [38,39]. The C100 appears to be a more potent neurotoxin than the A β [37]. It also exhibits neurotoxic actions *in vivo*, and transgenic mice overexpressing the peptide displays some neuropathologic features characteristic of the AD brain [40,41]. The processing of the APP is one critical event in the pathogenesis of AD; these processes can be modulated by different factors that include neurotransmitter receptors and activation of a multitude of second messenger systems. These can then provide therapeutic potentials for the action of anti-AD agents that may elicit certain pharmacologic action toward the increased secretion of soluble APPs and diminished production of the amyloidogenic A β peptide.

Tau protein—its role in AD progression

Tau (τ) is a major microtubule-associated protein that was almost simultaneously discovered in the United States and Europe [42–44]. It was first identified as a “factor essential for microtubule assembly” from purified tubulin. The τ protein is abundantly expressed in both the peripheral and the central nervous systems, [45] although low levels of τ protein are also present in the oligodendrocytes and astrocytes [46]. Tau undergoes a developmentally regulated phosphorylation having 38 phosphorylation sites [47]. The fetal brain as well as the pathologic brain has highly phosphorylated τ if compared with the normal adult brain [48]. The cDNA for τ was first isolated from the mouse brain expression library [49], then it was cloned from other species such as goat [50], chicken [51], bovine [52] and human [53,54].

The human τ gene is located on the human chromosome 17q21 and has at least 16 exons. To date, at least six different known major τ protein isoforms are brought about by messenger RNA (mRNA) alternative splicing [55] involving exons 2, 3, and 10, whereas exon 4A is only transcribed in the peripheral nervous system [56,57]. The expressed protein

is “the big τ ” isoform with approximate molecular weight of 100 kDa. The six isoforms consist of 352–441 amino acids with apparent molecular weight between approximately 37–46 kDa and pI values of 6.85–9.46 [58]. In some articles, the apparent molecular weights of the τ isoforms ranges from 60 kDa to 74 kDa [59]. The τ -isoform's structure becomes more complex due to various posttranslational modifications such as nitration, ubiquitination, glycosylation, phosphorylation, and truncation. Structurally, τ protein is hydrophilic and appears as a random coiled protein as analyzed by circular dichroism. The brain τ isoforms have two large domains: the projection domain, which comprises the amino acid terminal two-thirds of the molecule, and the microtubule-binding domain, which contains the carboxy-terminal third of the molecule. Furthermore, the projection domain can be divided into two regions: the amino terminal region with a high proportion of acidic residues and the proline-rich region. The microtubule binding domain can also be divided into the basic, true tubulin-binding region and the acidic carboxy-terminal.

AD is the best-known “tauopathy” because it is characterized by the aggregation of the abnormally phosphorylated τ protein. It is apparent that during normal development and as mentioned earlier, the τ protein undergoes various post-translational modifications. As a result, an increased amount of the modified τ has been associated with a large number of neurodegenerative disorders. In the process of aggregation to form paired helical filaments (PHFs), there should be a reduced affinity of the τ protein for the microtubules to accompany the release of τ into a soluble form. Dissociation of τ from the microtubule that may have been a result of the phosphorylation at some sites in the τ protein bring about microtubule destabilization. The resultant soluble τ protein becomes an ideal target for posttranslational modifications and may change directly or indirectly the conformation of the τ protein, thereby inducing τ dimerization in an antiparallel manner. Subsequently, the stable τ dimers then form τ oligomers and the aggregation process continues, therefore, constituting the subunits of filament called protomers. Coiling of two protomers with each other then forms PHFs. These PHFs then mature and assemble to form neurofibrillary tangles (NFTs). The NFTs are intraneuronal aggregates of the abnormally phosphorylated τ . NFTs correlate with the degree of dementia in AD. Currently, the molecular and cellular mechanisms that could explain the formation of τ lesions remain elusive. These lesions appeared to be controversial as to whether it could be considered a primary etiologic factor of the disease. Phosphorylation of τ is by far the most common and well-studied posttranslational modification associated with AD. It was observed that there is an obvious increased level of hyperphosphorylated τ in brains of AD patients that is three to four times higher than that found in normal adult brain [60,61] because normally a τ phosphoprotein contains one to three molecules of phosphate (PO_4^{3-}) per molecule of τ . Protein phosphorylation involves the addition of a phosphate group via esterification on the following amino acid residues: serine (S), threonine (T), and

tyrosine (Y). There are 85 putative phosphorylation sites on τ : 45 (53%) are on serines; 35 (41%) are on threonines; and 5 (6%) are on tyrosines. Phosphorylation is also modulated by various kinases and phosphatases. Therefore, hyperphosphorylation is a consequence of a dysregulation of the kinases and phosphatases. Currently there are more than 10 serine/threonine protein kinases known to phosphorylate τ protein *in vitro*. These kinases are categorized into two according to the motif specificity: proline-directed and nonproline-directed protein kinases. The proline-directed protein kinases include extracellular signal-regulated kinase 1/2 (erk-1/2), cell division cycle 2 kinase (cdc 2), cyclin-dependent kinase 2 (cdk 2), and the cdk 5. Although proline is not needed strictly in the phosphorylation of τ protein by Glycogen synthase-3 (GSK-3), still the presence of proline improves the phosphorylation efficiency. The nonproline-directed kinases are calcium- and calmodulin-dependent protein kinase II, PKA protein kinase (PKC), casein kinase I, casein kinase II and p¹¹⁰MARK.

The τ protein may also undergo prolyl isomerization, a reaction that allows the rearrangement of disulfide bonds in proteins. This involves the modification of the conformation of the protein of interest from *cis* to *trans* conformation and vice versa. The change in the conformation of the τ protein from *cis* to *trans* allows a shift of the peptide chain in space, allowing them to be more available for the action of phosphatases. Prolyl isomerization of τ can be achieved by the aid of the enzyme, peptidylprolyl isomerase Pin-1 (peptidylprolyl *cis/trans* isomerase NIMA interacting-1; NIMA: never in mitosis gene A) [62]. Pin-1 interacts with τ to prevent its pathologic accumulation [63]. Also, τ nitration proceeds by the addition of nitrogen dioxide on tyrosines of the molecule that occurs particularly at four sites Y18, [64] Y29, Y197, and Y394 [65]. This reaction was observed to be involved in the aggregation of the τ protein [66]. Polyamination is a reaction catalyzed by transglutaminases involving a glutamine residue as an acyl donor and a lysine as an acyl acceptor. The reaction results to the formation of γ -glutamyl- ϵ -lysine isopeptide bond and provokes protein cross-linking [67]. The primary donor site is Q424 and the principal acceptor is in the region between residues K163 and K240, [68] thereby linking τ polyamination to the NFT formation process. Oxidation can also occur on the τ protein that leads to the assembly of the paired helical filaments [69,70]; it occurs at amino acid residue C322 localized in the R3 domain of the microtubule-binding domain (MDB) region. The C322 is linked to the formation of tau lesions.

To date, it is still difficult to ascertain which among the previously described post-translational modifications is preferentially involved in τ pathology and AD. Most certainly, an orchestration among several τ modifications is required for τ aggregation into NFTs. Therefore, studying τ regulation at transcription and translation levels is of high significance in our understanding of the physiologic role of τ and its involvement in the development of AD. This is also particularly important in identifying compounds that are of therapeutic value for AD.

Curcuminoids and resveratrol on AD

Scientists around the globe have been trying for decades to treat and put an end to the pathologic symptoms of AD that deprive elderly persons of their intellect, but have had limited success. Despite these efforts, only four Food and Drug Administration-approved drugs are currently commercially available in the United States to treat AD pathology [71]. These drugs mainly target cholinergic functions associated with AD, leaving most other potential AD targets almost unaffected by the treatment. As a result, there is an urgent need for developing drugs based on multiple pathomechanisms of AD. The enormous diversity of functions in natural compounds extracted and isolated from either plant or marine sources may provide a new generation of drugs for AD therapy and management [72,73]. Plant-derived drugs are popular because the public believes that herbs are naturally safer than synthetic drugs. These beliefs may account for the sudden increase in the use of herbal treatment during the last decades [74]. This review also presents an update on the current researches undertaken on curcumin, demethoxycurcumin, and resveratrol regarding their potential for use in the treatment of AD.

Curcuminoids

Curcumin (Fig. 1) is a bright yellow powder that was first identified in 1910 [75]. This compound is also known as deferuloylmethane, natural yellow 3 and (1E,6E)-1,7-bis(4-hydroxy-3-methoxyphenyl)hepta-1,6-diene-3,5-dione. It has a formula weight equivalent to 368.38 g mole⁻¹. This substance was isolated from the plant *Curcuma longa* or turmeric that is commonly used for culinary purposes. This substance possesses several functional groups wherein the planar aromatic ring systems are attached to α,β -unsaturated carbonyl groups. The diketones generate stable enols that could be easily deprotonated to form enolates. The α,β -unsaturated group can undergo nucleophilic addition and Michael reaction. Curcumin has been shown to possess a wide range of pharmacologic activities, including anti-inflammatory [76,77], antioxidant [78], anticancer [79], wound healing [80], and antimicrobial effects. Curcumin has been shown to exhibit activity against various neurologic diseases, including AD [81], multiple sclerosis [82], Parkinson disease [83], epilepsy [84], cerebral injury [85], age-associated neurodegeneration [86], schizophrenia [87], spongiform encephalopathies (Creutzfeldt–Jakob disease) [88], neuropathic pain [89], and depression [90].

The oxidation process occurs mostly in the brain and it consumes 20% of the body's oxygen despite accounting for only 2% of the total body weight. With normal aging, the brain

accumulates metals ions such as iron (Fe), zinc (Zn), and copper (Cu). Thus, the brain requires copious amounts of antioxidants that control and prevent the damaging effects of reactive oxygen species generated via the Fenton reaction concerning redox-active metal-ion reduction and activation of molecular oxygen [91]. In AD, copper ions are also involved in the progression of the disease, and copper uptake is regulated by the APP. Copper has high affinity to both APP and A β ; Huang et al [92] recently reported that curcumin may play a dual role in protecting the rat cortical neuron against the Cu(II)-induced oxidative damage. Low dosages of curcumin (i.e. 1, 5, and 10 μ M) were able to depress oxidative stress levels exacerbated by Cu(II); however, high dosage (>10 μ M) of curcumin failed to decrease the Cu(II)-induced oxidative stress [92]. It was found that curcumin and its structural analogs, demethoxycurcumin and bisdemethoxycurcumin, can protect PC12 rat pheochromocytoma and normal human umbilical vein endothelial cells from A β -induced oxidative stress, and that these compounds were better antioxidants than α -tocopherol [93]. Zhang et al [94] in 2010 investigated the effect of curcumin on the metabolism of A β and APP in various cell lines including rat neuroblastoma cells over-expressing human APP₇₅₁. Curcumin significantly increased the levels of matured APP at low concentrations (5 and 10 μ M) and there was an observed decrease in the level of matured APP at 15 and 20 μ M. As a result, curcumin treatment markedly increased the ratio of matured APP and immature APP at lower concentrations and again the ratio was decreased significantly at higher concentrations of curcumin [94]. In a study by Ishrat et al [95], supplementation of curcumin on intracerebroventricular-streptozotocin infused rats significantly alleviated cognitive deficiencies resulted from intracerebroventricular-streptozotocin infusion. Curcumin further prevented oxidative damage to the hippocampus and cerebral cortex as ascertained through the following biomarkers: (1) increased 4-hydroxynonenal, malondialdehyde, thiobarbituric reactive substances, hydrogen peroxide, protein carbonyl and oxidized glutathione; and (2) diminished amounts of reduced glutathione and enzymes glutathione peroxidase and glutathione reductase [95]. The potential reversal of the intracerebroventricular-streptozotocin-induced alterations in cognitive behavioral and biochemical parameters suggests that addition of curcumin as part of our diet may help reduce the incidence of age-related sporadic dementia of Alzheimer type.

In 2005, Yang et al [96] showed that curcumin could inhibit aggregated as well as disaggregated A β 40. In this regard, curcumin was seen to be a better A β 40 aggregation inhibitor than ibuprofen or naproxen. In this regard, studies by Yanagisawa et al [97] on derivatization of curcumin aiming to increase the anti-A β aggregation activity of curcumin revealed that the curcumin derivatives have high levels of binding to A β aggregates and not the A β monomers. In the same study, the enol form was more predominant among the curcumin derivatives that interacted with the A β aggregates. This may be because compounds may only be able to bind with A β aggregates. In addition, the compound needs to be coplanar




Fig. 1. Structural formula of curcumin.

and have double-bond conformation with certain optimal carbon chains also known as the Goldilocks model [98]. The curcumin derivatives used in the study by Yanasigawa et al [97] conformed with the Goldilocks model for the compound, and maintained the coplanarity and the presence of the extended double bond conjugation between the two phenyl rings which gave the strong binding affinity of curcumin to the A β aggregates. The microtubule-associated protein 2 (MAP2) is a neuronal cytoskeletal component that plays an important role in the preservation of cellular architecture and internal organization, with certain involvement in the determination of cellular shape, in cell division, and cellular processes [99]. Changes in the architectural framework of the cytoskeleton bring forth malfunctions of the nerve cells in brain tissues of patients with AD. In a study by Xiao et al [100], it was reported that chemically synthesized curcumin and its derivatives can increase the viability of human neuroblastoma SK-N-SH cells after A β ₁₋₄₂ insult. Treatment of SH-N-SK cells with curcumin and its derivatives resulted in an increase expression of the MAP2. The upregulation of the MAP2 protein may be associated with the reorganization of the cytoskeleton during the degeneration of neuronal cells in AD and this implied that curcumin may have played a pivotal role in improving abnormal neuronal cell morphology by being able to restore normal physiology within the cell. In addition, the molecular structure of curcumin and its derivatives may also be responsible for their neuroprotective function. The important role of β -diketone moiety, 4-hydroxyl group on the benzene ring might be a key group of resistance to A β neurotoxicity, whereas 3-methoxy could indirectly affect the beneficial effect of curcumin derivatives related to suppressing 4-hydroxyl [101]. It was also described that the hydrogen atom donation from the β -diketone part of the lipid alkyl or the lipid peroxy radical is the potentially more important antioxidant action of curcumin [93]. Recently, solid-state nuclear magnetic resonance studies demonstrated that curcumin can interact with the 12th and 17th to 21st amino acid residues of the A β ₄₂ fibrils. Furthermore, based on the clear cross-peaks between carbons adjacent to methoxy and/or hydroxy groups revealed by C¹³ to C¹³ 2D solid-state nuclear magnetic resonance experiments, the presence of the functional groups on the aromatic rings plays an important role in the anti-A β aggregative property of curcumin [102]. In a study conducted by Liu et al [103], curcumin exhibited the most potent activity in the restraint of the production of A β ₄₂ in swap HEK293 cells followed by demethoxycurcumin and bisdemethoxycurcumin. Their group also found that curcumin but not other curcuminoids was able to decrease the APP protein expression, suggesting that the anti-amyloidogenic property of curcumin can be attributed to its effect on the APP metabolism.

Demethoxycurcumin (Fig. 2) is a structural analog of curcumin less one methoxy group and is also isolated from the plant *C. longa*. It comes as orange powder with a formula weight equal to 338.35 g mole⁻¹. Its IUPAC name is (1E,6E)-1-(4-hydroxy3-methoxyphenyl)-7-(4-hydroxyphenyl) hepta-1,6-diene-3,5-dione and may commonly be referred to as desmethoxycurcumin, monodemethoxycurcumin, or




Fig. 2. Structural formula of demethoxycurcumin.

parahydroxycurcumin. Several articles have been published associating members of the curcuminoid family with AD. Listed herein are some of the results gathered by researchers implicating demethoxycurcumin with AD. β -Amyloid (A β) induced oxidative stress is a well-established pathway of neuronal cell death in AD. The curcuminoids curcumin, demethoxycurcumin, and bisdemethoxycurcumin, were observed to protect PC12 rat pheochromocytoma and normal human umbilical vein endothelial cells from A β ₄₂ insult, as measured by MTT assay. The ED₅₀ values of curcumin, demethoxycurcumin, and bisdemethoxycurcumin toward PC12 and human umbilical vein endothelial cells were 7.1 \pm 0.3, 4.7 \pm 0.1, 3.5 \pm 0.2 μ g/mL and 6.8 \pm 0.4, 4.2 \pm 0.3, and 3.0 \pm 0.3 μ g/mL, respectively. These compounds were found to be better antioxidants than α -tocopherol as determined by DPPH radical trapping experiment. α -Tocopherol did not protect the cells from A β (1–42) insult even at >50 μ g/mL concentration. The results suggested that these compounds protected the cells from A β ₄₂ insult through antioxidant pathway [104]. Ahmed and Gilani [105] in 2009 reported on their conducted experiments on curcuminoids including demethoxycurcumin for anti-acetylcholinesterase (AChE) activity *in vitro* and *ex vivo*. It was observed that, *in vitro*, demethoxycurcumin has a marginal effect on AChE inhibition compared with bisdemethoxycurcumin and curcumin has the lowest activity. *Ex vivo*, both demethoxycurcumin and bisdemethoxycurcumin were able to show anti-AChE activity [105]. Investigation on the effects of curcuminoid mixture and individual constituents on spatial learning and memory in an A β peptide-infused rat model of AD and on the expression of post-synaptic density protein-95 (PSD-95), synaptophysin, and Ca²⁺/calmodulin-dependent protein kinase IV (camkIV) were carried out. Curcuminoid mixture exhibited a memory-enhancing effect in rats displaying AD-like neuronal loss marginally at 30 mg/kg, whereas individual components were effective at 3–30 mg/kg dosages. A short period of treatment with test compounds showed that the curcuminoid mixture and bisdemethoxycurcumin increased PSD-95 expression in the hippocampus at 3–30 mg/kg. The maximum effect was observed at a low dose of 3 mg/kg with respective values of 470.5 and 587.9%. However, after a longer treatment time, both demethoxycurcumin and curcumin also increased PSD-95 level up to 331.7 and 226.2%, respectively, at 30 mg/kg. Meanwhile, their effect on synaptophysin in the hippocampus after the longer duration treatment was studied. The curcuminoid mixture and all three individual components increased synaptophysin expression. Among these, demethoxycurcumin was the most effective, showing a 350.1% increase ($p < 0.01$) at 30 mg/kg compared with the neurotoxin group. The compounds' effect

on camkIV expression in the hippocampus was also determined. Only demethoxycurcumin increased camkIV levels to 421.2% at a concentration of 30 mg/kg. These compounds salvaged PSD-95, synaptophysin, and camkIV at the expression levels in the hippocampus in the rat AD model, suggesting multiple target sites giving out the potential of curcuminoids in enhancing the spatial memory and disease progression in AD. Recently, the effect of a curcuminoid mixture and its individual components on inflammatory and apoptotic genes expression in AD using an A β + ibotenic acid-infused rat model was explored. Five days after treatment with demethoxycurcumin, the amounts of hippocampal IL-1 β were decreased to 118.54 ± 47.48 and $136.67 \pm 31.96\%$, respectively, at 30 mg/kg and 10 mg/kg, compared with the amyloid treated group ($373.99 \pm 15.28\%$). Also, 5 days treatment with the curcuminoid mixture and demethoxycurcumin effectively decreased the levels of glial fibrillary acidic protein, a specific marker for astrocytes hyperactivity, often elevated in inflammatory conditions and AD, in the hippocampus. The curcuminoid mixture and bisdemethoxycurcumin effectively decreased caspase-3 level in the hippocampus after 20 days of treatment, where bisdemethoxycurcumin showed a maximal rescuing effect that is $92.35 \pm 3.07\%$ at a concentration of 3 mg/kg. Furthermore, the curcuminoid mixture also at 30 mg/kg decreased hippocampal Fas ligand levels to $70.56 \pm 3.36\%$ after a 5-day treatment and $19.01 \pm 2.03\%$ after 20 days. Measurement of the amount Fas receptor, demethoxycurcumin decreased the levels after 5 days of treatment, with all three doses showing a maximum effect ($189.76 \pm 15.01\%$) at 10 mg/kg. Individual compound was efficient 20 days posttreatment in plummeting Fas receptor levels in the hippocampus. This study demonstrated the important outcomes of curcuminoids administration on gene expressions.

Resveratrol

Resveratrol (Fig. 3) was first reported in 1939 by a Japanese researcher, Dr. Michio Takaoka, after it was extracted, purified, and isolated from a poisonous but medicinal herb, *Veratrum album* var. *grandiflorum* [106]. Resveratrol is a stilbenoid, a derivative of stilbene, that is also known as 3,4,5'-trihydroxystilbene. It has a molecular weight equivalent to $228.24 \text{ g mole}^{-1}$. This compound appears as white powder with slight yellow tinge. It exists in nature as two geometric isomers, *cis*- and *trans*-resveratrol. The *trans*-resveratrol is the more biologically active isomer. However, exposure to ultraviolet light of the *trans*-resveratrol leads to its conversion to its inactive form, *cis*-resveratrol [107]. In grapes,




Fig. 3. Structural formula of resveratrol.

trans-resveratrol is a phytoalexin produced against the growth of fungal pathogens such as *Botrytis cinerea* [108]. Also, most of this compound is found in the skin and seed of the grape's fruit (*Vitis vinifera*) [109]. An epidemiologic association between the common red wine and a low occurrence of cardiovascular diseases led to the beginning of what was referred to as the "French Paradox", coined by Serge Renaud, a scientist from Bordeaux University, despite the saturated fat-rich diet consumed by the French people. Furthermore, resveratrol attracted greater attention when its presence in red wine was suggested as an explanation for the cardioprotective effects of wine. Wine and grape extract have been shown to reduce platelet aggregation, uphold vasorelaxation, prevent atherosclerosis, decrease lipid peroxidation, and ameliorate serum cholesterol and triglyceride concentrations [110].

Because resveratrol has shown unprecedented promise for the development of new drugs, intense efforts are being made to understand its use in promoting healthy aging, longevity, and neuronal protection. Reports were made telling the significance of resveratrol as a neuroprotective agent. To date, it is still not clear how resveratrol may be essential in neuronal protection. Scientists currently still have to prove whether resveratrol takes its action through the traditionally known antioxidant properties that could be more effective than the cytoplasmic signal transduction pathways, or whether it may act through direct modulation of neuronal functions. Studies strongly implicate sirtuins in neuronal protection, and have also shown that calorie-restriction types and calorie-restriction mimetics can elevate sirtuin production in the human brain. As mentioned previously, resveratrol can act as a calorie-restriction mimetic, making it a good neuroprotective candidate compound acting via the sirtuin pathway [110]. Sirtuins, particularly SIRT-1, may play an important role in protecting neurons from the devastating effects of reactive oxygen species, peroxides, nitric oxide, β -amyloid peptides, and other intracellular and extracellular insults that could be present in the brain with AD. Resveratrol-induced SIRT-1 overexpression has been found to deacetylate and suppress the activity of p53 in neurons, thus preventing apoptotic death of neurons. In addition to this finding, the apoptotic activities of the FOXO proteins were also inhibited by resveratrol-induced SIRT-1 expression [111]. FOXOs has functional similarities and takes part in the cross-talk with p53. In another study, it was observed that resveratrol-induced SIRT-1 inhibited the signaling pathway of the nuclear factor kappa B (NF- κ B) in microglia and astrocytes resulting in the protection against A β -induced toxicity [112]. Soluble forms of A β can accumulate in the endothelial cells of the blood vessels of the brain with AD that ultimately inhibits the endothelial nitric oxide synthase (eNOS) activity. This phenomenon later causes toxic effects on nitric oxide functions such as Ca²⁺ dyshomeostasis [113]. This study suggests that in brains with AD, resveratrol may increase eNOS, modulate the expression of SIRT-1, and affect other cellular metabolic functions. Resveratrol also protects both nerves and blood vessels against A β insults. Moreover, amyloid peptides trigger the activation by acting with a number of Toll-like receptors (TLRs) including TLR4.

It was shown by Capiralla et al [114] that resveratrol can prevent the activation of murine RAW 264.7 macrophages (mouse leukaemic monocyte macrophage cell line) and the microglial BV-2 cells treated with the TLR4 ligand, lipopolysaccharide. The activation may be due to the reason that resveratrol may have acted upstream during the activation cascades interfering in the TLR4 oligomerization after the stimulation of the receptors. In this same study, resveratrol was demonstrated to hinder the proinflammatory events as consequences of the presence of the fibrillar A β on macrophages by effectively inhibiting the effect of A β on the phosphorylation of I κ B, activation of STAT1 and STAT3, and on the secretion of tumor necrosis factor- α and interleukin-6 [114].

In transgenic APP mice experiments, excessive and persistent reactive oxygen species production caused by the interaction of A β with A β -induced alcohol dehydrogenase may result in oxidative damage to mitochondrial and cellular biomolecules, resulting in a shut-down of mitochondrial energy production [115]. Finally, resveratrol had long been considered as a powerful antioxidant, and evidence has indicated the potential of resveratrol as a neuro-protecting agent. In PC12 cells, resveratrol was able to protect cells from A β_{25-35} -induced toxicity; it was also able to attenuate apoptotic cell death, reduce changes in mitochondrial potential and inhibited accumulation of reactive oxygen species [116]. Another important finding by Marambaud et al [117] regarding the neuroprotective role of resveratrol is its ability to interact with the ubiquitin proteasome system (UPS) by affecting the proteasome-mediated degradation of A β through a mechanism that does not directly increase the activity of the proteasome. The UPS is one of the main cellular, protein quality control mechanisms that enzymatically labels, transports, and eventually degrades the misprocessed and misfolded proteins [118]. It also controls all the metabolic reactions of the components of APP. Unfortunately it has been found that accumulation of A β causes the reduction of the UPS. Recently, it has been demonstrated that resveratrol was cytoprotective in human neuroblastoma cells by exerting its role not through the inhibition of β -amyloid aggregation but rather on its scavenging abilities from the toxic effect of the accumulation of A β and A β -metal complexes (A β -Fe, A β -Cu, A β -Zn, and A β -Al).

Resveratrol at a concentration of 15 μ M acted as a reactive oxygen species scavenger against those generated by A β -Fe, A β -Cu, A β -Zn complexes, thereby reducing their toxicity; however, resveratrol is not sufficient to fully block the toxicity induced by metal complexes [119]. Furthermore, it was previously reported in an *in vivo* experiment that 2% resveratrol did not alter the holoAPP, α C-terminal (C83), and β -terminal fragments (C99) when it was given *ad libitum* to transgenic APP695 mice, Tg19959 [120].

The poor bioavailability of resveratrol limits its potential for use in the treatment of AD because (1) polyphenols such as resveratrol undergo extensive biochemical reactions including oxidation, reduction, hydrolyses, glucuronic acid, sulfate, and methyl conjugation immediately after ingestion, and (2) there is a considerable person-to-person variability in drug absorption and metabolic process [109]. However, it is presumed that the circulating form of resveratrol in the human body, thought

to modulate numerous cellular and biochemical effects, is predominantly the modified metabolite and not the aglycone. This finding is true because the total recovery of glucuronic acid and sulfates conjugated with resveratrol in human urine and feces were about 71–98% through oral routes and 54–91% via the intravenous route [121].

Conclusion

AD is an incapacitating disease that deprives elderly individuals of their will and intellect. Apart from the diverse biologic qualities such as anti-inflammatory, antioxidant, anticancer, wound-healing enhancer, antimicrobial, and antiaging shown by curcumin, demethoxycurcumin, and resveratrol, these compounds had also found their niche in the anti-AD research. Researchers considered curcumin, demethoxycurcumin and resveratrol as possible anti-AD compounds due to their ability to prevent the aggregation of amyloid- β peptides. For many of the pharmacologic activities including the anti-AD activities that have been reported on curcumin, demethoxycurcumin, and resveratrol, clinical data are very limited. Clinical efficiency and potential toxicity of these compounds in larger trials requires further evaluation before recommendation regarding their use for AD treatment.

References

- [1] Whitehouse PJ, Price DL, Clark AW, Coyle JT, De Long MR. Alzheimer's disease: evidence for selective loss of cholinergic neurons in the nucleus basalis. *Ann Neurol* 1981;10:122–6.
- [2] Bouchard RW, Rossor MN. Typical clinical features. In: Gauthier S, editor. *Clinical diagnosis and management of Alzheimer's disease*. 3rd ed.; 2006. p. 39–52.
- [3] Kwok JBJ, Taddei K, Hallupp M, Fisher C, Brooks WS, Broe GA, et al. Two novel (M233T and [rho]278T) presenilin-1 mutations in early-onset Alzheimer's disease pedigrees and preliminary evidence for association of presenilin-1 mutations with a novel phenotype. *Neuroreport* 1997;8:1537–42.
- [4] Möller HJ, Graeber MB. The case described by Alois Alzheimer in 1911. *Eur Arch Psychiatry Clin Neurosci* 1998;248:111–22.
- [5] Alzheimer A. Über eine eigenartige Erkrankung der Hirnrinde. *Allgemeine Zeitschrift fuer Psychiatrie und psychisch-gerichtliche Medizin* 1907;64:146–8.
- [6] Small DH, Cappai R. Alois Alzheimer and Alzheimer's disease: a centennial perspective. *J Neurochem* 2006;99:708–10.
- [7] Ferri CP, Prince M, Brayne C, Brodaty H, Fratiglioni L, Ganguli M, et al. Global prevalence of dementia: a Delphi consensus study. *Lancet* 2005;366:2112–7.
- [8] WHO Alzheimer's disease: The Brain Killer. http://www.searo.who.int/en/Section1174/Section1199/Section1567/Section1823_8066.htm [14.02.12].
- [9] Dias A, Patel V. Closing the treatment gap for dementia in India. *Indian J Psych* 2009;51:93–7.
- [10] Lewis J, Dickson DW, Lin WL, Chisholm L, Corral A, Jones G, et al. Enhanced neurofibrillary degeneration in transgenic mice expressing mutant tau and APP. *Science* 2001;293:1487–91.
- [11] Kang J, Lemaire HG, Unterbeck A, Salbaum JM, Masters CL, Grzeschik KH, et al. The precursor of Alzheimer's disease amyloid A4 protein resembles a cell-surface receptor. *Nature* 1987;325:733–6.
- [12] Yankner BA, Duffy LK, Kirschner DA. Neurotrophic and neurotoxic effects of amyloid beta protein: reversal by tachykinin neuropeptides. *Science* 1990;250:279–82.

- [13] Hartley DM, Walsh DM, Ye CP, Diehl T, Vasquez S, Vassilev PM, et al. Protofibrillar intermediates of amyloid beta-protein induce acute electrophysiological changes and progressive neurotoxicity in cortical neurons. *J Neurosci* 1999;19:8876–84.
- [14] Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, et al. Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A* 1998;95:6448–53.
- [15] Walsh DM, Klyubin I, Fadeeva JV, Cullen WK, Anwyl R, Wolfe MS, et al. Naturally secreted oligomers of amyloid beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature* 2002;416:535–9.
- [16] Ono K, Condon MM, Teplow DB. Structure-neurotoxicity relationships of amyloid β -protein oligomers. *Proc Natl Acad Sci* 2009;106:14745–50.
- [17] Podlisny MB, Ostaszewski BL, Squazzo SL, Koo EH, Rydel RE, Teplow DB, et al. Aggregation of secreted amyloid beta-protein into sodium dodecyl sulfate-stable oligomers in cell culture. *J Biol Chem* 1995;270:9564–70.
- [18] Walsh DM, Tseng BP, Rydel RE, Podlisny MB, Selkoe DJ. The oligomerization of amyloid beta-protein begins intracellularly in cells derived from human brain. *Biochemistry* 2000;39:10831–9.
- [19] Turner RS, Suzuki N, Chyung AS, Younkin SG, Lee VM. Amyloids beta40 and beta42 are generated intracellularly in cultured human neurons and their secretion increases with maturation. *J Biol Chem* 1996;271:8966–70.
- [20] Coughlan CM, Breen KC. Factors influencing the processing and function of the amyloid β precursor protein—a potential therapeutic target in Alzheimer's disease? *Pharmacol Therapeut* 2000;86:111–44.
- [21] Jarrett JT, Lansbury Jr PT. Seeding “one-dimensional crystallization” of amyloid: a pathogenic mechanism in Alzheimer's disease and scrapie? *Cell* 1993;73:1055–8.
- [22] Lansbury Jr PT. Consequences of the molecular mechanism of amyloid formation for the understanding of the pathogenesis of Alzheimer's disease and the development of therapeutic strategies. *Arzneimittelforschung* 1995;45:432–4.
- [23] Suzuki N, Cheung TT, Cai XD, Odaka A, Otvos Jr L, Eckman C, et al. An increased percentage of long amyloid beta protein secreted by familial amyloid beta protein precursor (beta APP717) mutants. *Science* 1994;264:1336–40.
- [24] Scheuner D, Eckman C, Jensen M, Song X, Citron M, Suzuki N, et al. Secreted amyloid beta-protein similar to that in the senile plaques of Alzheimer's disease is increased in vivo by the presenilin 1 and 2 and APP mutations linked to familial Alzheimer's disease. *Nat Med* 1996;2:864–70.
- [25] Goate A, Chartier-Harlin MC, Mullan M, Brown J, Crawford F, Fidani L, et al. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 1991;349:704–6.
- [26] Breen KC. APP-collagen interaction is mediated by a heparin bridge mechanism. *Mol Chem Neuropathol* 1992;16:109–21.
- [27] Kibbey MC, Jucker M, Weeks BS, Neve RL, Van Nostrand WE, Kleinman HK. Beta-Amyloid precursor protein binds to the neurite-promoting IKVAV site of laminin. *Proc Natl Acad Sci U S A* 1993;90:10150–3.
- [28] Narindrasorasak S, Altman RA, Gonzalez-DeWhitt P, Greenberg BD, Kisilevsky R. An interaction between basement membrane and Alzheimer amyloid precursor proteins suggests a role in the pathogenesis of Alzheimer's disease. *Lab Invest* 1995;72:272–82.
- [29] Coulson EJ, Barrett GL, Storey E, Bartlett PF, Beyreuther K, Masters CL. Down-regulation of the amyloid protein precursor of Alzheimer's disease by antisense oligonucleotides reduces neuronal adhesion to specific substrata. *Brain Res* 1997;770:72–80.
- [30] Multhaup G. Identification and regulation of the high affinity binding site of the Alzheimer's disease amyloid protein precursor (APP) to glycosaminoglycans. *Biochimie* 1994;76:304–11.
- [31] Sisodia SS, Gallagher M. A role for the beta-amyloid precursor protein in memory? *Proc Natl Acad Sci U S A* 1998;95:12074–6.
- [32] Moser EI, Krobot KA, Moser MB, Morris RG. Impaired spatial learning after saturation of long-term potentiation. *Science* 1998;281:2038–42.
- [33] Mattson MP, Cheng B, Culwell AR, Esch FS, Lieberburg I, Rydel RE. Evidence for excitoprotective and intraneuronal calcium-regulating roles for secreted forms of the beta-amyloid precursor protein. *Neuron* 1993;10:243–54.
- [34] Goodman Y, Mattson MP. Secreted forms of beta-amyloid precursor protein protect hippocampal neurons against amyloid beta-peptide-induced oxidative injury. *Exp Neurol* 1994;128:1–12.
- [35] Jin LW, Ninomiya H, Roch JM, Schubert D, Masliah E, Otero DA, et al. Peptides containing the RERMS sequence of amyloid beta/A4 protein precursor bind cell surface and promote neurite extension. *J Neurosci* 1994;14:5461–70.
- [36] Pike CJ, Overman MJ, Cotman CW. Amino-terminal deletions enhance aggregation of beta-amyloid peptides in vitro. *J Biol Chem* 1995;270:23895–8.
- [37] Kim SH, Suh YH. Neurotoxicity of a carboxyl-terminal fragment of the Alzheimer's amyloid precursor protein. *J Neurochem* 1996;67:1172–82.
- [38] Kim HS, Park CH, Suh YH. C-terminal fragment of amyloid precursor protein inhibits calcium uptake into rat brain microsomes by Mg^{2+} - Ca^{2+} ATPase. *Neuroreport* 1998;9:3875–9.
- [39] Kim HS, Lee JH, Suh YH. C-terminal fragment of Alzheimer's amyloid precursor protein inhibits sodium/calcium exchanger activity in SK-N-SH cell. *Neuroreport* 1999;10:113–6.
- [40] Kammesheidt A, Boyce FM, Spanoyannis AF, Cummings BJ, Ortegón M, Cotman C, et al. Deposition of beta/A4 immunoreactivity and neuronal pathology in transgenic mice expressing the carboxyl-terminal fragment of the Alzheimer amyloid precursor in the brain. *Proc Natl Acad Sci U S A* 1992;89:10857–61.
- [41] Pascale A, Bhagavan S, Nelson TJ, Neve RL, McPhie DL, Etcheberrigaray R. Enhanced BK-induced calcium responsiveness in PC12 cells expressing the C100 fragment of the amyloid precursor protein. *Mol Brain Res* 1999;72:205–13.
- [42] Cleveland DW, Hwo SY, Kirschner MW. Purification of tau, a microtubule-associated protein that induces assembly of microtubules from purified tubulin. *J Mol Biol* 1977;116:207–25.
- [43] Cleveland DW, Hwo SY, Kirschner MW. Physical and chemical properties of purified tau factor and the role of tau in microtubule assembly. *J Mol Biol* 1977;116:227–47.
- [44] Fellous A, Francon J, Lennon AM, Nunez J. Microtubule assembly in vitro. Purification of assembly-promoting factors. *Eur J Biochem* 1977;78:167–74.
- [45] Binder LI, Frankfurter A, Rebhun LI. The distribution of tau in the mammalian central nervous system. *J Cell Biol* 1985;101:1371–8.
- [46] LoPresti P, Szuchet S, Papasozomenos SC, Zinkowski RP, Binder LI. Functional implications for the microtubule-associated protein tau: localization in oligodendrocytes. *Proc Natl Acad Sci U S A* 1995;92:10369–73.
- [47] Pittman AM, Fung H-C, de Silva R. Untangling the tau gene association with neurodegenerative disorders. *Hum Mol Gen* 2006;15:R188–95.
- [48] Kanamaru K, Takio K, Miura R, Titani K, Ihara Y. Fetal-type phosphorylation of the tau in paired helical filaments. *J Neurochem* 1992;58:1667–75.
- [49] Lee G, Cowan N, Kirschner M. The primary structure and heterogeneity of tau protein from mouse brain. *Science* 1988;239:285–8.
- [50] Nelson PT, Stefansson K, Gulcher J, Saper CB. Molecular evolution of tau protein: implications for Alzheimer's disease. *J Neurochem* 1996;67:1622–32.
- [51] Yoshida H, Goedert M. Molecular cloning and functional characterization of chicken brain tau: isoforms with up to five tandem repeats. *Biochemistry* 2002;41:15203–11.
- [52] Himmler A, Drechsel D, Kirschner MW, Martin Jr DW. Tau consists of a set of proteins with repeated C-terminal microtubule-binding domains and variable N-terminal domains. *Mol Cell Biol* 1989;9:1381–8.
- [53] Goedert M, Spillantini MG, Cairns NJ, Crowther RA. Tau proteins of Alzheimer paired helical filaments: abnormal phosphorylation of all six brain isoforms. *Neuron* 1992;8:159–68.
- [54] Goedert M, Spillantini MG, Potier MC, Ulrich J, Crowther RA. Cloning and sequencing of the cDNA encoding an isoform of microtubule-

- associated protein tau containing four tandem repeats: differential expression of tau protein mRNAs in human brain. *EMBO J* 1989;8:393–9.
- [55] Neve RL, Harris P, Kosik KS, Kurnit DM, Donlon TA. Identification of cDNA clones for the human microtubule-associated protein tau and chromosomal localization of the genes for tau and microtubule-associated protein 2. *Brain Res* 1986;387:271–80.
 - [56] Andreadis A, Broderick JA, Kosik KS. Relative exon affinities and suboptimal splice site signals lead to non-equivalence of two cassette exons. *Nucleic Acids Res* 1995;23:585–93.
 - [57] Buee L, Bussiere T, Buee-Scherrer V, Delacourte A, Hof PR. Tau protein isoforms, phosphorylation and role in neurodegenerative disorders. *Brain Res Brain Res Rev* 2000;33:95–130.
 - [58] Wang J-Z, Liu F. Microtubule-associated protein tau in development, degeneration and protection of neurons. *Prog Neurobiol* 2008;85:148–75.
 - [59] Martin L, Latypova X, Terro F. Post-translational modifications of tau protein: implications for Alzheimer's disease. *Neurochemistry Int* 2011;58:458–71.
 - [60] Khatoon S, Grundke-Iqbal I, Iqbal K. Levels of normal and abnormally phosphorylated tau in different cellular and regional compartments of Alzheimer disease and control brains. *FEBS Lett* 1994;351:80–4.
 - [61] Khatoon S, Grundke-Iqbal I, Iqbal K. Brain levels of microtubule-associated protein tau are elevated in Alzheimer's disease: a radio-immuno-slot-blot assay for nanograms of the protein. *J Neurochem* 1992;59:750–3.
 - [62] Yaffe MB, Schutkowski M, Shen M, Zhou XZ, Stukenberg PT, Rahfeld JU, et al. Sequence-specific and phosphorylation-dependent proline isomerization: a potential mitotic regulatory mechanism. *Science* 1997;278:1957–60.
 - [63] Yotsumoto K, Saito T, Asada A, Oikawa T, Kimura T, Uchida C, et al. Effect of Pin1 or microtubule binding on dephosphorylation of FTDP-17 mutant Tau. *J Biol Chem* 2009;284:16840–7.
 - [64] Reyes JF, Reynolds MR, Horowitz PM, Fu Y, Guillozet-Bongaarts AL, Berry R, et al. A possible link between astrocyte activation and tau nitration in Alzheimer's disease. *Neurobiol Dis* 2008;31:198–208.
 - [65] Reynolds MR, Berry RW, Binder LI. Site-specific nitration differentially influences tau assembly in vitro. *Biochemistry* 2005;44:13997–4009.
 - [66] Horiguchi T, Uryu K, Giasson BI, Ischiropoulos H, Lightfoot R, Bellmann C, et al. Nitration of tau protein is linked to neurodegeneration in tauopathies. *Am J Pathol* 2003;163:1021–31.
 - [67] Wang DS, Dickson DW, Malter JS. Tissue transglutaminase, protein cross-linking and Alzheimer's disease: review and views. *Int J Clin Exp Pathol* 2008;1:5–18.
 - [68] Murthy SN, Wilson JH, Lukas TJ, Kuret J, Lorand L. Cross-linking sites of the human tau protein, probed by reactions with human transglutaminase. *J Neurochem* 1998;71:2607–14.
 - [69] Schweers O, Mandelkow EM, Biernat J, Mandelkow E. Oxidation of cysteine-322 in the repeat domain of microtubule-associated protein tau controls the in vitro assembly of paired helical filaments. *Proc Natl Acad Sci U S A* 1995;92:8463–7.
 - [70] Landino LM, Skreslet TE, Alston JA. Cysteine oxidation of tau and microtubule-associated protein-2 by peroxynitrite: modulation of microtubule assembly kinetics by the thioredoxin reductase system. *J Biol Chem* 2004;279:35101–5.
 - [71] Allain H, Bentue-Ferrer D, Tribut O, Gauthier S, Michel BF, Drieu-La Rochelle C. Alzheimer's disease: the pharmacological pathway. *Fundam Clin Pharmacol* 2003;17:419–28.
 - [72] Stafford GI, Pedersen ME, van Staden J, Jager AK. Review on plants with CNS-effects used in traditional South African medicine against mental diseases. *J Ethnopharmacol* 2008;119:513–37.
 - [73] Howes MJ, Houghton PJ. Plants used in Chinese and Indian traditional medicine for improvement of memory and cognitive function. *Pharmacol Biochem Behav* 2003;75:513–27.
 - [74] Raskin I, Ribnicky DM, Komarnytsky S, Ilic N, Poulev A, Borisjuk N, et al. Plants and human health in the twenty-first century. *Trends Biotechnol* 2002;20:522–31.
 - [75] Miłobędzka J, Kostanecki S, Lampe V. Zur Kenntnis des Curcumins. *Berichte der deutschen chemischen Gesellschaft* 1910;43:2163–70.
 - [76] Srivastava R, Dikshit M, Srimal RC, Dhawan BN. Anti-thrombotic effect of curcumin. *Thromb Res* 1985;40:413–7.
 - [77] Abe Y, Hashimoto SHU, Horie T. Curcumin inhibition inflammatory cytokine production by human peripheral blood monocytes and alveolar macrophages. *Pharmacol Res* 1999;39:41–7.
 - [78] Sharma OP. Antioxidant activity of curcumin and related compounds. *Biochem Pharmacol* 1976;25:1811–2.
 - [79] Adams BK, Ferstl EM, Davis MC, Herold M, Kurtkaya S, Camalier RF, et al. Synthesis and biological evaluation of novel curcumin analogs as anti-cancer and anti-angiogenesis agents. *Bioorg Med Chem* 2004;12:3871–83.
 - [80] Sidhu GS, Singh AK, Thaloor D, Banaudha KK, Patnaik GK, Srimal RC, et al. Enhancement of wound healing by curcumin in animals. *Wound Repair Regen* 1998;6:167–77.
 - [81] Lim GP, Chu T, Yang F, Beech W, Frautschy SA, Cole GM. The curry spice curcumin reduces oxidative damage and amyloid pathology in an Alzheimer transgenic mouse. *J Neurosci* 2001;21:8370–7.
 - [82] Natarajan C, Bright JJ. Curcumin inhibits experimental allergic encephalomyelitis by blocking IL-12 signaling through Janus kinase-STAT pathway in T lymphocytes. *J Immunol* 2002;168:6506–13.
 - [83] Zbarsky V, Datla KP, Parkar S, Rai DK, Aruoma OI, Dexter DT. Neuroprotective properties of the natural phenolic antioxidants curcumin and naringenin but not quercetin and fisetin in a 6-OHDA model of Parkinson's disease. *Free Radic Res* 2005;39:1119–25.
 - [84] Sumanont Y, Murakami Y, Tohda M, Vajragupta O, Watanabe H, Matsumoto K. Prevention of kainic acid-induced changes in nitric oxide level and neuronal cell damage in the rat hippocampus by manganese complexes of curcumin and diacetylcurcumin. *Life Sci* 2006;78:1884–91.
 - [85] Ghoneim AI, Abdel-Naim AB, Khalifa AE, El-Denshary ES. Protective effects of curcumin against ischaemia/reperfusion insult in rat forebrain. *Pharmacol Res* 2002;46:273–9.
 - [86] Calabrese V, Scapagnini G, Colombrita C, Ravagna A, Pennisi G, Giuffrida Stella AM, et al. Redox regulation of heat shock protein expression in aging and neurodegenerative disorders associated with oxidative stress: a nutritional approach. *Amino Acids* 2003;25:437–44.
 - [87] Bishnoi M, Chopra K, Kulkarni SK. Protective effect of Curcumin, the active principle of turmeric (*Curcuma longa*) in haloperidol-induced orofacial dyskinesia and associated behavioural, biochemical and neurochemical changes in rat brain. *Pharmacol Biochem Behav* 2008;88:511–22.
 - [88] Hafner-Bratkovic I, Gaspersic J, Smid LM, Bresjanac M, Jerala R. Curcumin binds to the alpha-helical intermediate and to the amyloid form of prion protein – a new mechanism for the inhibition of PrP(Sc) accumulation. *J Neurochem* 2008;104:1553–64.
 - [89] Sharma S, Kulkarni SK, Agrewala JN, Chopra K. Curcumin attenuates thermal hyperalgesia in a diabetic mouse model of neuropathic pain. *Eur J Pharmacol* 2006;536:256–61.
 - [90] Xu Y, Ku BS, Yao HY, Lin YH, Ma X, Zhang YH, et al. The effects of curcumin on depressive-like behaviors in mice. *Eur J Pharmacol* 2005;518:40–6.
 - [91] Smith DG, Cappai R, Barnham KJ. The redox chemistry of the Alzheimer's disease amyloid beta peptide. *Biochim Biophys Acta* 2007;1768:1976–90.
 - [92] Huang H-C, Lin C-J, Liu W-J, Jiang Z- F. Dual effects of curcumin on neuronal oxidative stress in the presence of Cu(II). *Food Chem Toxicol* 2011;49:1578–83.
 - [93] Kim DS, Park SY, Kim JK. Curcuminoids from *Curcuma longa* L. (Zingiberaceae) that protect PC12 rat pheochromocytoma and normal human umbilical vein endothelial cells from betaA(1-42) insult. *Neurosci Lett* 2001;303:57–61.
 - [94] Zhang C, Browne A, Child D, Tanzi RE. Curcumin decreases amyloid-beta peptide levels by attenuating the maturation of amyloid-beta precursor protein. *J Biol Chem* 2010;285:28472–80.
 - [95] Ishrat T, Hoda MN, Khan MB, Yousuf S, Ahmad M, Khan MM, et al. Amelioration of cognitive deficits and neurodegeneration by curcumin in rat model of sporadic dementia of Alzheimer's type (SDAT). *Eur Neuropsychopharmacol* 2009;19:636–47.

- [96] Yang F, Lim GP, Begum AN, Ubeda OJ, Simmons MR, Ambegaokar SS, et al. Curcumin inhibits formation of amyloid-beta oligomers and fibrils, binds plaques and reduces amyloid *in vivo*. *J Biol Chem* 2005;280:5892–901.
- [97] Yanagisawa D, Shirai N, Amatsubo T, Taguchi H, Hirao K, Urushitani M, et al. Relationship between the tautomeric structures of curcumin derivatives and their A β -binding activities in the context of therapies for Alzheimer's disease. *Biomaterials* 2010;31:4179–85.
- [98] Reinke AA, Gestwicki JE. Structure-activity relationships of amyloid beta-aggregation inhibitors based on curcumin: influence of linker length and flexibility. *Chem Biol Drug Des* 2007;0:206–15.
- [99] Johnson GV, Jope RS. The role of microtubule-associated protein 2 (MAP-2) in neuronal growth, plasticity, and degeneration. *J Neurosci Res* 1992;33:505–12.
- [100] Xiao Z, Lin L, Liu Z, Ji F, Shao W, Wang M, et al. Potential therapeutic effects of curcumin: relationship to microtubule associated proteins 2 in Ab 1-2 insults. *Brain Res* 2010;1361:115–23.
- [101] Masuda T, Hidaka K, Shinohara A, Maekawa T, Takeda Y, Yamaguchi H. Chemical studies on antioxidant mechanism of curcuminoid: analysis of radical reaction products from curcumin. *J Agric Food Chem* 1999;47:71–7.
- [102] Masuda Y, Fukuchi M, Yatagawa T, Tada M, Takeda K, Irie K, et al. Solid-state NMR analysis of interaction sites of curcumin and 42-residue amyloid β -protein fibrils. *Bioorg Med Chem* 2011;19:5967–74.
- [103] Liu H, Li Z, Qiu D, Gu Q, Lei Q, Mao L. The inhibitory effects of different curcuminoids on β -amyloid protein, β -amyloid precursor protein and β -site amyloid precursor protein cleaving enzyme 1 in swAPP HEK293 cells. *Neurosci Lett* 2010;485:83–8.
- [104] Kim DSHL, Park S-Y, Kim J- Y. Curcuminoids from *Curcuma longa* L. (Zingiberaceae) that protect PC12 rat pheochromocytoma and normal human umbilical vein endothelial cells from β A(1-42) insult. *Neurosci Lett* 2001;303:57–61.
- [105] Ahmed T, Gilani A- H. Inhibitory effect of curcuminoids on acetylcholinesterase activity and attenuation of scopolamine-induced amnesia may explain medicinal use of turmeric in Alzheimer's disease. *Pharmacol Biochem Behav* 2009;91:554–9.
- [106] Takaoka M. Resveratrol, a new phenolic compound from *Veratrum grandiflorum*. *Nippon Kagaku Kaishi* 1939;60:1090–100.
- [107] Lamuela-Raventos RM, Romero-Perez AI, Waterhouse AL, de la Torre-Boronat MC. Direct HPLC analysis of cis- and trans-resveratrol and piceid isomers in Spanish red *Vitis vinifera* wines. *J Agric Food Chem* 1995;43:281–3.
- [108] Favaron F, Odorizzi S, Pais da Cunha AT, Sella L. The role of grape polyphenols on trans-resveratrol activity against *Botrytis cinerea* and of fungal laccase on the solubility of putative grape PR proteins. *J Plant Pathol* 2009;91:579–88.
- [109] Anekonda TS. Resveratrol – a boon for treating Alzheimer's disease? *Brain Res Rev* 2006;52:316–26.
- [110] Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the in vivo evidence. *Nat Rev Drug Discov* 2006;5:493–506.
- [111] Anekonda TS, Reddy PH. Neuronal protection by sirtuins in Alzheimer's disease. *J Neurochem* 2006;96:305–13.
- [112] Chen J, Zhou Y, Mueller-Steiner S, Chen LF, Kwon H, Yi S, et al. SIRT1 protects against microglia-dependent amyloid-beta toxicity through inhibiting NF-kappaB signaling. *J Biol Chem* 2005;280:40364–74.
- [113] Gentile MT, Vecchione C, Maffei A, Aretini A, Marino G, Poulet R, et al. Mechanisms of soluble beta-amyloid impairment of endothelial function. *J Biol Chem* 2004;279:48135–42.
- [114] Capiralla H, Vingetdeux V, Zhao H, Sankowski R, Al-Abed Y, Davies P, et al. Resveratrol mitigates lipopolysaccharide- and A β -mediated microglial inflammation by inhibiting the TLR4/NF- κ B/STAT signaling cascade. *J Neurochem* 2012;120:461–72.
- [115] Reddy PH, Beal MF. Are mitochondria critical in the pathogenesis of Alzheimer's disease? *Brain Res Brain Res Rev* 2005;49:618–32.
- [116] Jang JH, Surh YJ. Protective effect of resveratrol on beta-amyloid-induced oxidative PC12 cell death. *Free Radic Biol Med* 2003;34:1100–10.
- [117] Marambaud P, Zhao H, Davies P. Resveratrol promotes clearance of Alzheimer's disease amyloid-beta peptides. *J Biol Chem* 2005;280:37377–82.
- [118] de Vrij FM, Fischer DF, van Leeuwen FW, Hol EM. Protein quality control in Alzheimer's disease by the ubiquitin proteasome system. *Prog Neurobiol* 2004;74:249–70.
- [119] Granzotto A, Zatta P. Resveratrol acts not through anti-aggregative pathways but mainly via its scavenging properties against Abeta and Abeta-metal complexes toxicity. *PLoS One* 2011;6:e21565.
- [120] Karuppagounder SS, Pinto JT, Xu H, Chen H-L, Beal MF, Gibson GE. Dietary supplementation with resveratrol reduces plaque pathology in a transgenic model of Alzheimer's disease. *Neurochem Int* 2009;54:111–8.
- [121] Vitaglione P, Sforza S, Galaverna G, Ghidini C, Caporaso N, Vescovi PP, et al. Bioavailability of trans-resveratrol from red wine in humans. *Mol Nutr Food Res* 2005;49:495–504.