

Original Article

Bisphenol A at environmentally relevant doses induces cyclooxygenase-2 expression and promotes invasion of human mesenchymal stem cells derived from uterine myoma tissue

Kai-Hung Wang^{a,b,c}, An-Pei Kao^d, Chia-Cheng Chang^e, Ta-Chin Lin^{a,b}, Tsung-Cheng Kuo^{a,b,*}^aDepartment of Obstetrics and Gynecology, Kuo General Hospital, Tainan, Taiwan^bCenter for Reproductive Medicine, Kuo General Hospital, Tainan, Taiwan^cDepartment of Laboratory Medicine, Kuo General Hospital, Tainan, Taiwan^dDepartment of Obstetrics and Gynecology, Center for Reproductive Medicine and Sciences, Taipei Medical University Hospital, Taipei Medical University, Taipei, Taiwan^eDepartment of Pediatrics and Human Development, Michigan State University, East Lansing, MI 48824, USA

Accepted 17 October 2012

Abstract

Objective: Uterine myoma is the most common benign reproductive tract tumor in women. Despite its high prevalence, the exact pathogenesis of these benign tumors remains unknown. Toward understanding the pathogenic mechanism of these tumors, we attempted to isolate human uterine myoma mesenchymal stem cells (hUM-MSCs), which may be the target cells for tumorigenesis. Furthermore, we tested the response of these hUM-MSCs to the environmental endocrine disruptor, bisphenol A (BPA), which may mimic the action of estrogen in hormone-sensitive organs such as the uterus.

Materials and Methods: The hUM-MSC lines were clonally derived from uterine myoma tissue using the MSU-1 medium supplemented with *N*-acetyl-L-cysteine and L-ascorbic acid-2-phosphate. These hUM-MSCs were characterized by reverse transcription polymerase chain reaction (RT-PCR) analysis for the expression of mesenchymal stem cell (MSC) surface markers (e.g., CD90 and CD105) and the transcription factor Oct-4. The proliferation potential was measured by the cumulative population doubling level and the colony-forming efficiency.

Results: Putative hUM-MSC lines expressed *CD90*, *CD105*, and the stem cell marker gene, *Oct-4*. The cells were capable of differentiating into adipocytes, osteoblasts, and chondrocytes. Bisphenol A treatment of these hUM-MSCs enhanced cell proliferation and colony-forming efficiency in a dose-responsive manner. At an environmentally relevant concentration (10^{-8} M), BPA moreover induced *cyclooxygenase-2* (*COX-2*) gene expression and promoted cell migration and invasiveness.

Conclusion: The hUM-MSC cell lines can be isolated from uterine myoma tissues. Bisphenol A could enhance cell proliferation and colony-forming efficiency, induce *COX-2* gene expression, and promote migration and invasion of hUM-MSCs. The results imply that BPA has a detrimental effect on female health by promoting uterine tumorigenesis.

Copyright © 2013, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

Keywords: bisphenol A; cell invasion; cyclooxygenase-2; mesenchymal stem cells; uterine myoma

Introduction

Stem cells are undifferentiated cells with the capacity for unlimited or prolonged self-renewal. They are able to give rise

* Corresponding author. Department of Obstetrics and Gynecology, Kuo General Hospital, Number 22, Section 2, Minsheng Road, Tainan City, Taiwan.
E-mail address: tkuo@kgh.com.tw (T.-C. Kuo).

to differentiated cells. In recent years, tumor stem cells have been reported for many different tumors [1–8]. Many tumor stem cell markers are also markers of normal stem cells in different tissues [9–14], a major evidence supporting the hypothesis that stem cells are the tumor-initiating target cells [15,16].

A uterine myoma (i.e., a uterine fibroid) is a benign tumor of the female reproductive tract. Despite the high prevalence

of uterine myoma, its underlying pathogenesis is largely unknown [17]. One possible mechanism for uterine myoma formation is the dysregulation of stem cell activity. Evidence for the presence of uterine stem cells has been previously reviewed [18] and reported, and include eutopic and ectopic endometrial tissue [19], endometrial carcinoma [20], myometrium tissue [21], and myoma [22].

Mesenchymal stem cells (MSCs) have been defined by the International Society for Cellular Therapy [23] as possessing three characteristics, namely: (1) they are plastic-adherent; (2) they express CD90 and CD105, but not CD34 and CD45; and (3) they are able to differentiate into osteoblasts, adipocytes, and chondroblasts *in vitro*. The most common source of MSCs is the bone marrow [24]. Mesenchymal stem cells have been derived from various tissues such as adipose tissue [25], lipoma [26], cord blood [27], skeletal muscle, dermis [28], synovial membrane [29], placenta [30], amniotic fluid [31], eutopic and ectopic endometrial tissues [19], and peripheral blood [32].

Environmental endocrine disruptors (EEDs) include many environmental estrogens or xenoestrogens that mimic the action of natural estrogen hormone in the body. Bisphenol A (BPA), one of the most important EEDs, is widely used as a material for the production of epoxy resins and polycarbonate plastics. It has been detected in many human tissue compartments such as serum, follicular fluid, amniotic fluid [33,34], urine [35], breast milk [36], saliva [37], and adipose tissue [38] at concentrations ranging from 0.5 nM to 40 nM [34,39–43]. The consequence of BPA exposure in human reproduction remains controversial, although BPA exposure reportedly may be associated with recurrent miscarriages, endometrial hyperplasia, obesity, and polycystic ovarian syndrome (PCOS) [33,44–46].

In this study, we have developed a technique to isolate human uterine myoma tissue-derived mesenchymal stem cells (hUM-MSCs). This provides an opportunity to analyze the effects of BPA on hUM-MSCs.

Materials and methods

Primary culture cells derived from human uterine myoma tissue

Human uterine myoma tissue primary cells were isolated from five patients receiving surgery to treat uterine myoma in the Department of Obstetrics and Gynecology of Kuo General Hospital, Tainan, Taiwan. The use of these tissues has been approved by the Institutional Review Board (IRB). A small amount of uterine myoma tissue (approximately 5 grams) was minced by sterile scissors, and then subjected to overnight digestion at 37°C with collagenase Type I (1 mg/mL) in phenol red-free Dulbecco's modified Eagle's medium/F-12 medium (DMEM/F-12 medium) (5 mL medium per 1 g of tissue). The digested tissue was filtered through a sterile 100- μ m polyethylene mesh filter to remove undigested tissues, and again filtered through a 40- μ m cell strainer to remove epithelial cells. The primary cells in the filtrate were collected

by centrifugation and re-suspended in phenol red-free DMEM/F-12 medium supplemented with 10% fetal bovine serum (FBS). On the next day, unattached cells were removed by washing them with phosphate-buffered saline (PBS). The medium was changed every other day, and the cells were allowed to grow until near confluence. After 1 week, the primary cell cultures were trypsinized and stored in liquid nitrogen or subcultured for further experiments.

Isolation and culture of hUM-MSCs

To isolate hUM-MSCs, the primary cell cultures were diluted and seeded in a 96-well plate at a density of one cell per well. All wells were examined under the microscope for the number of attached cells. Wells without cells or with more than one cell were marked and disregarded. The medium was changed to the basic MSU-1 (i.e., phenol red-free) medium [47,48] with 5% FBS, 2 mM *N*-acetyl-L-cysteine (NAC) and 0.2 mM L-ascorbic acid-2-phosphate (Asc-2P). After incubation for 14 days, proliferating cells clonally derived from one single cell were trypsinized and cultured in a 60-mm Petri dish. The medium was renewed every other day and the cells were allowed to grow until near confluence. Three hUM-MSC cultures—hUM-MSC1, hUM-MSC2 and hUM-MSC3—were eventually obtained by this procedure. Early passage cells (i.e., passage 2) were used in subsequent experiments.

Cumulative population doubling level

The cumulative population doubling level (cpdl) in continual subcultures and growth from a known number of cells (1×10^5) were calculated to determine the proliferation potential of hUM-MSCs. The cpdl of each subculture was calculated from the cell count by using the following equation:

$$\ln(N_f/N_i)/\ln 2$$

in which N_i and N_f are the initial and final cell numbers, respectively, and \ln is the natural logarithm [19,49].

Cell proliferation assay

To examine the self-renewal capacity, the hUM-MSCs were treated for 7 days with various concentrations of 17- β -estradiol (E2), BPA (1×10^{-4} M, 1×10^{-5} M, 1×10^{-6} M, 1×10^{-7} M, 1×10^{-8} M, 1×10^{-9} M, and 1×10^{-10} M dissolved in dimethyl sulfoxide [DMSO]) or treated with the solvent vehicle DMSO (0.1%) as the negative control. The growth rates of the cell cultures were calculated by dividing the cpdl by the number of days that elapsed during the culture [19,47]. At least three independent experiments were performed for each study.

Colony formation assay

To test the effect of BPA on colony-forming ability, the hUM-MSCs cells were plated at a density of 100 viable cells per 10-cm dish in a medium without or with BPA (1×10^{-4} M, 1×10^{-5} M, 1×10^{-6} M, 1×10^{-7} M, 1×10^{-8} M, 1×10^{-9} M, and

1×10^{-10} M). After 21 days, the colonies were stained with 0.5% crystal violet stain. The colonies on the plates were photographed and counted to compare the colony-forming efficiency [19,47]. At least three independent experiments were performed for each study.

Multilineage differentiation potential of hUM-MSCs

The hUM-MSCs cells were tested for their mesodermal multilineage differentiation potential (i.e., adipogenesis, osteogenesis, and chondrogenesis). Cells were first subcultured and incubated in modified Eagle's medium (i.e., the D-medium) [50] with 10 % FBS. The next day, differentiation induction was initiated by changing the medium with different supplementations [19,25,49].

Adipogenesis

The hUM-MSCs were plated at a seeding density of 5×10^4 cells/cm² in 35-mm Petri dishes. The next day, the cells were incubated in IDII medium for 2 days, and then in I medium for 1 day. The cycle of treatment (i.e., 3 days) was repeated four times. The IDII medium contains 3-isobutyl-1-methylxanthine (500 μ M), dexamethasone (1 μ M), indomethacin (1 μ M), and insulin (10 μ g/mL) in D-medium with 10% FBS. The I medium is the D-medium supplemented with 10% FBS and insulin (10 μ g/mL). Oil Red O staining (indicated by red stain) verified the development of lipid vacuoles within adipocytes after induction.

Osteogenesis

The hUM-MSCs were plated at a seeding density of 5×10^4 cells/cm² in 35-mm Petri dishes. On the following day, the cells were induced to differentiate in DAG medium for 2 weeks. The medium was changed once every 3 days. The DAG medium contains 0.1 mM dexamethasone, 50 μ M L-ascorbic acid-2-phosphate, and 10 mM β -glycerophosphate disodium in D-medium with 10% FBS. The formation of calcified extracellular matrix (ECM) by osteoblasts was visible and can be confirmed by von Kossa staining (indicated by black stain).

Chondrogenesis

Micromass cultures of cells (1×10^5 cells/10 μ L) were incubated and formed in 24-well plates for 2.5 hours, and then induced to differentiate in TAI medium for 2 weeks. The medium was changed once every 3 days. The TAI medium contained 10 ng/mL TGF- β 1, 50 μ M L-ascorbic acid-2-phosphate, and 6.25 μ g/mL insulin in D-medium with 10% FBS. The micromass was stained with Alcian blue for the presence of sulfated proteoglycan-rich matrix (indicated by blue stain).

Migration and invasion assay

Cells were inoculated into 24-well Falcon Migration inserts (1×10^4 cells/well) and Matrigel Invasion inserts (1×10^5

cells/well) (8- μ m pore size in both inserts). Inserts were placed into Falcon companion plates and incubated for 18 hours for migration and 24 hours for invasion. After incubation, the medium and cells were removed from the top chamber by using cotton swabs and PBS. The number of cells that had invaded or migrated to the underside of the membrane were counted and recorded as the average number of migrating or invading cells per plate in triplicate plates [19,47,48,51].

Reverse transcription-polymerase chain reaction

Total RNA was extracted from cells by using 1 mL TRIzol. The RNA (1 μ g) was transcribed into cDNA with a reverse transcription system kit, and cDNA was amplified with gene-specific primers in a polymerase chain reaction (PCR) machine. The primers included *Oct-4* (forward, 5'-GGCTTGGAGACCTCTCAGCCTG; reverse, 5'-TGCAGCAAGGGCCGAGCTTAC), *CD90* (forward, 5'-TCGCTCTCCTGCTAACAGTCT; reverse, 5'-CTCGTACTGGATGGGTGAAC), *CD105* (forward, 5'-TG TCTCACTTCATGCCTCCAGCT; reverse, 5'-AGGCTGTCCA TGTTGAGGCAGT), *COX-2* (forward, 5'-AGGGCCAGCTTTC ACCAAC; reverse, 5'-AAGGCGCAGTTTACGCTGTC), and β -actin (loading control) (forward, 5'-ATGATATCGCCGCGC TCGTCGTC; reverse, 5'-CGCTCGGCCGTGGTGGTGAA). The amplified products were separated on 2% agarose gels and stained with ethidium bromide.

Statistical analysis

The results show the representatives of at least three separate experiments. The significance of difference between treatments was assessed by the Mann-Whitney test of nonparametric statistics and was performed by using the SPSS for Windows 13.0 statistics program (SPSS Inc., Chicago, IL, USA). A *p* value <0.05 was considered significant. All statistical data are presented as the mean \pm the standard deviation.

Results

Morphology of the primary cells and hUM-MSCs

Primary cells and hUM-MSCs were isolated from human uterine myoma tissue, as described in the Materials and methods section. Five primary cell cultures and three clonally derived hUM-MSC cell lines—hUM-MSC1, hUM-MSC2 and hUM-MSC3—were obtained by this procedure. In the following experiments, either all three hUM-MSC cell lines or just the hUM-MSC2 cell line was used as the representative cells for some studies. The primary cells were morphologically flat and large (Fig. 1A), whereas the hUM-MSCs were morphologically mostly fibroblast-like, but with a few serpiginous cells (i.e., cells with two long tails), and tended to show contact-insensitive growth (i.e., piling up) (Fig. 1A). As observed previously [25], some MSCs may appear as serpiginous cells. These serpiginous cells may divide symmetrically to give rise to two serpiginous cells or may divide asymmetrically to give rise to one serpiginous cell and one cuboidal cell [19,25,49].




Fig. 1. Morphology and stem cell characteristics of hUM-MSCs. (A) The morphology of primary cells is flat and large, whereas the morphology of hUM-MSCs is fibroblastlike or serpiginous. (B) Reverse transcription-polymerase chain reaction (RT-PCR) analysis shows that, in contrast to primary cells, hUM-MSC2 cells express *Oct-4* mRNA. The hUM-MSC1 and hUM-MSC3 cells have similar results (data are not shown). (C) The cumulative population doubling level of hUM-MSC1, hUM-MSC2, and hUM-MSC3 cells reached 20.6, 23.9, and 19.7, respectively, in 48 days (6 passages). hUM-MSC = human uterine myoma mesenchymal stem cell.

Expression of the *Oct-4* gene in hUM-MSCs

In the literature, the transcription factor Oct-4 is a stem cell marker that is expressed in embryonic stem cells, in germ cells [52], in many adult human stem cells [53], and in cancer stem cells [47,48]. Oct-4 expression appears essential for maintaining an undifferentiated state and preventing the differentiation of stem cells [54]. Based on RT-PCR analysis in this study, the *Oct-4* gene is expressed in hUM-MSCs, but not in primary cells (Fig. 1B).

Proliferation potential of hUM-MSCs

Stem cells are characterized by an extensive ability for self-renewal. To determine the proliferation potential of hUM-MSCs, the cumulative population doubling level (cpdl) of three hUM-MSC lines were measured in six passages during 48 days of cell growth. The cpdl for hUM-MSC1, hUM-MSC2, hUM-MSC3 were 20.6, 23.9, and 19.7, respectively (Fig. 1C).

In vitro differentiation and expression of MSC genes of hUM-MSCs

The qualifying criteria for cells to be considered as MSCs are (1) the ability to differentiate into cells of multiple tissue lineages and (2) the expression of CD90 and CD105 [23]. To demonstrate that the hUM-MSCs are capable of multilineage differentiation, they were grown in D-medium with different

supplementations to induce differentiation. After induction with specific medium supplementations, as described in the Materials and methods section, the hUM-MSCs differentiated into adipocytes, osteoblasts, and chondroblasts, as indicated by positive staining with Oil Red O (for lipid droplets), with von Kossa (for calcified ECM), and with Alcian blue (for sulfated proteoglycan), respectively (Fig. 2A). By RT-PCR analysis, specific MSC markers (i.e. *CD90* and *CD105*), were expressed in the hUM-MSC2 (Fig. 2B). *CD90* and *CD105* were also expressed in hUM-MSC1 and hUM-MSC3 (data are not shown).

BPA enhances the growth rate and clonogenic ability of hUM-MSCs in a dose-responsive manner

The growth response of hUM-MSC2 to E2 and BPA concentrations ranging from 10^{-10} M to 10^{-4} M was analyzed. The results showed that E2 concentrations from 10^{-10} M to 10^{-5} M significantly increased cell growth, compared to the vehicle control. The optimum dose was from 10^{-8} M to 10^{-6} M. The BPA treatment (at a concentration of 10^{-8} M to 10^{-5} M) of hUM-MSC2 cells also significantly increased cell growth. Both chemicals inhibited cell growth at 10^{-4} M (Fig. 3A). In the clonogenic assay, E2 and BPA significantly increased the colony formation of hUM-MSC2 cells in a dose-responsive manner at a concentration range of 10^{-10} M to 10^{-6} M (Fig. 3B). In the tested dose range, the effects of E2 were more significant than the effects of BPA. A BPA concentration of 10^{-4} M inhibited clonogenic ability. The effective concentration of BPA at 10^{-8} M is considered an environmentally relevant dose and was used for subsequent experiments.

BPA at an environmentally relevant dose induced cyclooxygenase-2 (*COX-2*) gene expression in hUM-MSCs cell

In the medical literature, prolonged exposure of breast stem cells to E2 could enhance COX-2 expression [47]. Many




Fig. 2. In vitro differentiation and expression of MSC genes of hUM-MSCs. (A) The hUM-MSCs were induced to differentiate into the following mesenchymal lineages: adipocytes (stained red by using Oil Red O staining of lipid droplets); osteoblasts (stained black by using von Kossa staining of calcified extracellular matrix [ECM]); and chondroblasts (stained blue by using Alcian blue staining of sulfated proteoglycans). (B) Reverse transcription-polymerase chain reaction (RT-PCR) analysis indicated *CD90* and *CD105* mRNA expression in the hUM-MSC2 cell lines. BPA = bisphenol A; hUM-MSC = human uterine myoma mesenchymal stem cell.




Fig. 3. Bisphenol A (BPA) enhances the growth rate and clonogenic ability of hUM-MSCs. (A) The effect of 17- β -estradiol (E2) or BPA on cell proliferation in hUM-MSC2 cells. The cells treated with various concentrations of E2 or BPA for 7 days were subjected to the proliferation assay. The ratio of the value for the E2-treated or BPA-treated cells to the value for the control cells (i.e., DMSO-treated cells) was calculated. These are shown on the ordinate. Values are presented as the mean \pm standard deviation (SD) ($n = 3$, different cultures). (B) The hUM-MSC2 cells treated with various concentrations of E2, BPA, or DMSO (0.1%). After 21 days, colonies of cells were stained with 0.5% crystal violet. The values are presented as the mean \pm SD ($n = 3$, different cultures). BPA = bisphenol A; DMSO = dimethyl sulfoxide; E2 = 17- β -estradiol; hUM-MSC = human uterine myoma mesenchymal stem cell. * $p < 0.05$.

environmental endocrine disruptors may mimic the action of natural hormone estrogens such as E2 in the body. Therefore, we investigated the effect of BPA treatment on COX-2 expression in hUM-MSCs. After treatment with BPA (10^{-8} M) for two weeks, the COX-2 mRNA level in hUM-MSC2 cells was significantly increased, based on RT-PCR analysis (Fig. 4A). Similar results were obtained for hUM-MSC1 and hUM-MSC3 (data are not shown).

BPA at an environmentally relevant dose enhances the migration and invasion ability of hUM-MSCs cells

The effects of BPA to promote cell migration and invasion ability of hUM-MSCs were examined by using migration and invasion chamber assays. The results reveal that BPA

(10^{-8} M) treatment of hUM-MSC2 for two weeks significantly increased cell migration ability, compared to hUM-MSC2 without treatment (2.5% vs. 1.6%, respectively) (Fig. 4B). The same concentration of BPA similarly significantly enhanced the invasion ability of hUM-MSC2 (0.238% vs. 0.086% for the control) (Fig. 4B).

Discussion

This study demonstrated that MSCs can be isolated from human uterine myoma tissue by using the basic MSU-1 medium [47,48,55] supplemented with 5% FBS, 2 mM NAC, and 0.2 mM Asc-2P. The MSU-1 medium without NAC and Asc-2P was first used in developing human breast epithelial stem cells [55]. We previously used a keratinocyte medium supplemented with NAC and Asc-2P to develop MSCs from human and other mammalian adipose tissues [25,26,49,56,57]. The success of developing these MSCs using these media is ascribed to using NAC [58,59] and Asc-2p [60–62] antioxidants which maintain a cellular redox state that is essential for stem cell self-renewal [58].

The hUM-MSCs in this study possessed many MSC phenotypes. First, the cells showed a high proliferation potential (more than 20–24 cpdl for the three clonally derived hUM-MSC cell lines that were examined). Second, the cells were fibroblast-like or serpiginous with a morphology similar to adipose tissue-derived human MSCs [25], lipoma [26], and eutopic or ectopic endometrial tissue [19]. They showed contact-insensitive growth (i.e., piling up). Third, the hUM-MSCs expressed MSC markers (i.e., CD90 and CD105) and the stem cell marker Oct-4, which is expressed in many adult human and canine stem cells [53,56]. Fourth, and most importantly, the hUM-MSCs were capable of differentiating into three major mesodermal cell lineages: adipocytes, osteoblasts, and chondroblasts.

Using the hUM-MSCs, we tested the effect of BPA on gene expression and on phenotypic changes. The results show that BPA increased hUM-MSC proliferation and colony-forming




Fig. 4. BPA induces COX-2 gene expression and promotes the migration and invasion of hUM-MSCs. (A) Reverse transcription-polymerase chain reaction (RT-PCR) analysis shows that the COX-2 mRNA expression is significantly elevated in hUM-MSC2 cells treated with BPA (10^{-8} M) for 2 weeks, compared to the controls (i.e., without BPA treatment). (B) Based on migration and invasion chamber assay, hUM-MSC2 cells treated with BPA (10^{-8} M) for two weeks have significantly enhanced cell migration and invasion. The values are presented as the mean \pm standard deviation (SD) ($n = 3$, different cultures). BPA = bisphenol A; hUM-MSC = human uterine myoma mesenchymal stem cell. * $p < 0.05$.

efficiency in a dose-dependent manner. Furthermore, long-term BPA exposure at environmentally relevant doses induced *COX-2* gene expression and promoted the migration and invasion ability of hUM-MSCs. It has been reported previously that BPA could promote proliferation of MCF-7 cells [63], HeLa cells [64], OVCAR-3 cell line [65], neuroblastoma [66], and JKT-1 cells [67]. The growth-promoting effect of BPA on hUM-MSCs has not been reported previously. In the future, the results derived from experiments using hUM-MSC2 may be confirmed by using the other 2 hUM-MSC lines.

The elevated expression of *COX-2* could elicit many effects such as the promotion of tumor progression by inducing invasion and angiogenesis, the suppression of host immunity, the resistance to apoptosis, and the epithelial to mesenchymal transition (see references cited in Wang et al [47]). We previously reported that interleukin-1 beta could induce *COX-2* overexpression and promote tumor invasion [51]. The results of this study indicate that long-term exposure of hUM-MSCs to BPA at environmentally relevant doses may induce *COX-2* gene expression and may enhance invasion ability, which is consistent with previous reports [51,68].

In conclusion, we have developed a technique to isolate clonally derived hUM-MSC lines. We further demonstrated that BPA could enhance cell proliferation and colony-forming efficiency, induce *COX-2* gene expression, and promote migration and invasion ability of hUM-MSCs. The results provide evidence that BPA could promote uterine tumor growth. These hUM-MSC lines could be useful for detecting environmental endocrine disruptors or toxicants that promote tumorigenesis.

Acknowledgments

This work was supported by the Kuo General Hospital Research Fund (grant numbers 100-E-004, 100-E-005, and 100-E-025).

References

- [1] Bonnet D, Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med* 1997;3:730–7.
- [2] Singh SK, Hawkins C, Clarke ID, Squire JA, Bayani J, Hide T, et al. Identification of human brain tumour initiating cells. *Nature* 2004;432:396–401.
- [3] Kim CF, Jackson EL, Woolfenden AE, Lawrence S, Babar I, Vogel S, et al. Identification of bronchioalveolar stem cells in normal lung and lung cancer. *Cell* 2005;121:823–35.
- [4] Fang D, Nguyen TK, Leishear K, Finko R, Kulp AN, Hotz S, et al. A tumorigenic subpopulation with stem cell properties in melanomas. *Cancer Res* 2005;65:9328–37.
- [5] Li C, Heidt DG, Dalerba P, Burant CF, Zhang L, Adsay V, et al. Identification of pancreatic cancer stem cells. *Cancer Res* 2007;67:1030–7.
- [6] O'Brien CA, Pollett A, Gallinger S, Dick JE. A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature* 2007;445:106–10.
- [7] Collins AT, Berry PA, Hyde C, Stower MJ, Maitland NJ. Prospective identification of tumorigenic prostate cancer stem cells. *Cancer Res* 2005;65:10946–51.
- [8] Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA* 2003;100:3983–8.
- [9] Hope KJ, Jin L, Dick JE. Acute myeloid leukemia originates from a hierarchy of leukemic stem cell classes that differ in self-renewal capacity. *Nat Immunol* 2004;5:738–43.
- [10] Singh SK, Clarke ID, Terasaki M, Bonn VE, Hawkins C, Squire J, et al. Identification of a cancer stem cell in human brain tumors. *Cancer Res* 2003;63:5821–8.
- [11] Sell S. Alpha-fetoprotein, stem cells and cancer: how study of the production of alpha-fetoprotein during chemical hepatocarcinogenesis led to reaffirmation of the stem cell theory of cancer. *Tumour Biol* 2008;29:161–80.
- [12] Herrera MB, Bruno S, Buttiglieri S, Tetta C, Gatti S, Deregibus MC, et al. Isolation and characterization of a stem cell population from adult human liver. *Stem Cells* 2006;24:2840–50.
- [13] McClanahan T, Koseoglu S, Smith K, Grein J, Gustafson E, Black S, et al. Identification of overexpression of orphan G protein-coupled receptor GPR49 in human colon and ovarian primary tumors. *Cancer Biol Ther* 2006;5:419–26.
- [14] Becker L, Huang Q, Mashimo H. Immunostaining of Lgr5, an intestinal stem cell marker, in normal and premalignant human gastrointestinal tissue. *ScientificWorldJournal* 2008;8:1168–76.
- [15] Chang CC. Recent translational research: stem cells as the roots of breast cancer. *Breast Cancer Res* 2006;8:103.
- [16] Lin T, Meng L, Li Y, Tsai RY. Tumor-initiating function of nucleostemin-enriched mammary tumor cells. *Cancer Res* 2010;70:9444–52.
- [17] Stewart EA. Uterine fibroids. *Lancet* 2001;357:293–8.
- [18] Maruyama T, Masuda H, Ono M, Kajitani T, Yoshimura Y. Human uterine stem/progenitor cells: their possible role in uterine physiology and pathology. *Reproduction* 2010;140:11–22.
- [19] Kao AP, Wang KH, Chang CC, Lee JN, Long CY, Chen HS, et al. Comparative study of human eutopic and ectopic endometrial mesenchymal stem cells and the development of an in vivo endometriotic invasion model. *Fertil Steril* 2011;95:1308–15.
- [20] Hubbard SA, Friel AM, Kumar B, Zhang L, Rueda BR, Gargett CE. Evidence for cancer stem cells in human endometrial carcinoma. *Cancer Res* 2009;69:8241–8.
- [21] Ono M, Maruyama T, Masuda H, Kajitani T, Nagashima T, Arase T, et al. Side population in human uterine myometrium displays phenotypic and functional characteristics of myometrial stem cells. *Proc Natl Acad Sci USA* 2007;104:18700–5.
- [22] Chang HL, Senaratne TN, Zhang L, Szotek PP, Stewart E, Dombkowski D, et al. Uterine leiomyomas exhibit fewer stem/progenitor cell characteristics when compared with corresponding normal myometrium. *Reprod Sci* 2010;17:158–67.
- [23] Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini F, Krause D, et al. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 2006;8:315–7.
- [24] Conget PA, Minguell JJ. Phenotypical and functional properties of human bone marrow mesenchymal progenitor cells. *J Cell Physiol* 1999;181:67–73.
- [25] Lin TM, Tsai JL, Lin SD, Lai CS, Chang CC. Accelerated growth and prolonged lifespan of adipose tissue-derived human mesenchymal stem cells in a medium using reduced calcium and antioxidants. *Stem Cells Dev* 2005;14:92–102.
- [26] Lin TM, Chang HW, Wang KH, Kao AP, Chang CC, Wen CH, et al. Isolation and identification of mesenchymal stem cells from human lipoma tissue. *Biochem Biophys Res Comm* 2007;361:883–9.
- [27] Lee OK, Kuo TK, Chen WM, Lee KD, Hsieh SL, Chen TH. Isolation of multipotent mesenchymal stem cells from umbilical cord blood. *Blood* 2004;103:1669–75.
- [28] Young HE, Steele TA, Bray RA, Hudson J, Floyd JA, Hawkins K, et al. Human reserve pluripotent mesenchymal stem cells are present in the connective tissues of skeletal muscle and dermis derived from fetal, adult, and geriatric donors. *Anat Rec* 2001;264:51–62.

- [29] De Bari C, Dell'Accio F, Tylzanowski P, Luyten FP. Multipotent mesenchymal stem cells from adult human synovial membrane. *Arthritis Rheum* 2001;44:1928–42.
- [30] Yen BL, Huang HI, Chien CC, Jui HY, Ko BS, Yao M, et al. Isolation of multipotent cells from human term placenta. *Stem Cells* 2005;23:3–9.
- [31] Tsai MS, Lee JL, Chang YJ, Hwang SM. Isolation of human multipotent mesenchymal stem cells from second-trimester amniotic fluid using a novel two-stage culture protocol. *Hum Reprod* 2004;19:1450–6.
- [32] Kassis I, Zangi L, Rivkin R, Leviansky L, Samuel S, Marx G, et al. Isolation of mesenchymal stem cells from G-CSF-mobilized human peripheral blood using fibrin microbeads. *Bone Marrow Transplant* 2006;37:967–76.
- [33] Takeuchi T, Tsutsumi O. Serum bisphenol A concentrations showed gender differences, possibly linked to androgen levels. *Biochem Biophys Res Commun* 2002;291:76–8.
- [34] Ikezuki Y, Tsutsumi O, Takai Y, Kamei Y, Taketani Y. Determination of bisphenol A concentrations in human biological fluids reveals significant early prenatal exposure. *Hum Reprod* 2002;17:2839–41.
- [35] Ouchi K, Watanabe S. Measurement of bisphenol A in human urine using liquid chromatography with multi-channel coulometric electrochemical detection. *J Chromatography. B, Anal Tech in Biomed and Life Sci* 2002;780:365–70.
- [36] Ye X, Kuklennyk Z, Needham LL, Calafat AM. Measuring environmental phenols and chlorinated organic chemicals in breast milk using automated on-line column-switching-high performance liquid chromatography-isotope dilution tandem mass spectrometry. *J Chromatography. B, Anal Tech in Biomed and Life Sci* 2006;831:110–5.
- [37] Sasaki N, Okuda K, Kato T, Kakishima H, Okuma H, Abe K, et al. Salivary bisphenol-A levels detected by ELISA after restoration with composite resin. *J Mater Sci Mater Med* 2005;16:297–300.
- [38] Fernandez MF, Arrebola JP, Taoufik J, Navalón A, Ballesteros O, Pulgar R, et al. Bisphenol-A and chlorinated derivatives in adipose tissue of women. *Reprod Toxicol* 2007;24:259–64.
- [39] Calafat AM, Kuklennyk Z, Reidy JA, Caudill SP, Ekong J, Needham LL. Urinary concentrations of bisphenol A and 4-nonylphenol in a human reference population. *Environ Health Perspect* 2005;113:391–5.
- [40] Braun JM, Kalkbrenner AE, Calafat AM, Bernert JT, Ye X, Silva MJ, et al. Variability and predictors of urinary bisphenol A concentrations during pregnancy. *Environ Health Perspect* 2011;119:131–7.
- [41] Sun Y, Irie M, Kishikawa N, Wada M, Kuroda N, Nakashima K. Determination of bisphenol A in human breast milk by HPLC with column-switching and fluorescence detection. *Biomed Chromatogr* 2004;18:501–7.
- [42] Vandenberg LN, Hauser R, Marcus M, Olea N, Welshons WV. Human exposure to bisphenol A (BPA). *Reprod Toxicol* 2007;24:139–77.
- [43] Welshons WV, Nagel SC, vom Saal FS. Large effects from small exposures. III. Endocrine mechanisms mediating effects of bisphenol A at levels of human exposure. *Endocrinology* 2006;147:S56–69.
- [44] Takeuchi T, Tsutsumi O, Ikezuki Y, Takai Y, Taketani Y. Positive relationship between androgen and the endocrine disruptor, bisphenol A, in normal women and women with ovarian dysfunction. *Endocr J* 2004;51:165–9.
- [45] Sugiura-Ogasawara M, Ozaki Y, Sonta S, Makino T, Suzumori K. Exposure to bisphenol A is associated with recurrent miscarriage. *Hum Reprod* 2005;20:2325–9.
- [46] Yang M, Park MS, Lee HS. Endocrine disrupting chemicals: human exposure and health risks. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev* 2006;24:183–224.
- [47] Wang KH, Kao AP, Chang CC, Lee JN, Chai CY, Hou MF, et al. Modulation of tumorigenesis and oestrogen receptor- α expression by cell culture conditions in a stem cell-derived breast epithelial cell line. *Biol Cell* 2010;102:159–72.
- [48] Wang KH, Kao AP, Chang CC, Lee JN, Hou MF, Long CY, et al. Increasing CD44⁺/CD24⁺ tumor stem cells, and upregulation of COX-2 and HDAC6, as major functions of HER2 in breast tumorigenesis. *Mol Cancer* 2010;9:288.
- [49] Wang KH, Kao AP, Chen HW, Wang FY, Chang CH, Chang CC, et al. Optimizing proliferation and characterization of multipotent stem cells from porcine adipose tissue. *Biotech Applied Biochem* 2008;51:159–66.
- [50] Chang CC, Boezi JA, Warren ST, Sabourin CL, Liu PK, Glatzer L, et al. Isolation and characterization of a UV-sensitive hypermutable aphidicolin-resistant Chinese hamster cell line. *Somatic Cell Genet* 1981;7:235–53.
- [51] Kao AP, Wang KH, Long CY, Chai CY, Tsai CF, Hsieh TH, et al. Interleukin-1 β induces cyclooxygenase-2 expression and promotes the invasive ability of human mesenchymal stem cells derived from ovarian endometrioma. *Fertil Steril* 2011;96:678–84.
- [52] Scholer HR, Ruppert S, Suzuki N, Chowdhury K, Gruss P. New type of POU domain in germ line-specific protein Oct-4. *Nature* 1990;344:435–9.
- [53] Tai MH, Chang CC, Kiupel M, Webster JD, Olson LK, Trosko JE. Oct4 expression in adult human stem cells: evidence in support of the stem cell theory of carcinogenesis. *Carcinogenesis* 2005;26:495–502.
- [54] Pesce M, Scholer HR. Oct-4: gatekeeper in the beginnings of mammalian development. *Stem Cells* 2001;19:271–8.
- [55] Kao CY, Nomata K, Oakley CS, Welsch CW, Chang CC. Two types of normal human breast epithelial cells derived from reduction mammaplasty: phenotypic characterization and response to SV40 transfection. *Carcinogenesis* 1995;16:531–8.
- [56] Neupane M, Chang CC, Kiupel M, Yuzbasiyan-Gurkan V. Optimization of growth and differentiation of canine adipose-derived mesenchymal stem cells. *Tissue Eng* 2008;14:1007–15.
- [57] Wang KH, Kao AP, Singh S, Yu SL, Kao LP, Tsai ZY, et al. Comparative expression profiles of mRNAs and microRNAs among human mesenchymal stem cells derived from breast, face, and abdominal adipose tissues. *Kaohsiung J Med Sci* 2010;26:113–22.
- [58] Smith J, Ladi E, Mayer-Proschel M, Noble M. Redox state is a central modulator of the balance between self-renewal and differentiation in a dividing glial precursor cell. *Proc Natl Acad Sci USA* 2000;97:10032–7.
- [59] De Flora S, Izzotti A, D'Agostini F, Balansky RM. Mechanisms of N-acetylcysteine in the prevention of DNA damage and cancer, with special reference to smoking-related end-points. *Carcinogenesis* 2001;22:999–1013.
- [60] Ivanov VO, Ivanova SV, Niedzwiecki A. Ascorbate affects proliferation of guinea-pig vascular smooth muscle cells by direct and extracellular matrix-mediated effects. *J Mol Cell Cardiol* 1997;29:3293–303.
- [61] Hata R, Senoo H. L-ascorbic acid 2-phosphate stimulates collagen accumulation, cell proliferation, and formation of a three-dimensional tissue-like substance by skin fibroblasts. *J Cell Physiol* 1989;138:8–16.
- [62] Chepda T, Cadau M, Girin P, Frey J, Chamson A. Monitoring of ascorbate at a constant rate in cell culture: effect on cell growth. *In Vitro Cell Dev Biol Anim* 2001;37:26–30.
- [63] Ricupito A, Del Pozzo G, Diano N, Grano V, Portaccio M, Marino M, et al. Effect of bisphenol A with or without enzyme treatment on the proliferation and viability of MCF-7 cells. *Environ Int* 2009;35:21–6.
- [64] Bolli A, Galluzzo P, Ascenzi P, Del Pozzo G, Manco I, Vietri MT, et al. Laccase treatment impairs bisphenol A-induced cancer cell proliferation affecting estrogen receptor α -dependent rapid signals. *IUBMB Life* 2008;60:843–52.
- [65] Ptak A, Wróbel A, Gregoraszczuk EL. Effect of bisphenol-A on the expression of selected genes involved in cell cycle and apoptosis in the OVCAR-3 cell line. *Toxicol Lett* 2011;202:30–5.
- [66] Zhu H, Xiao X, Zheng J, Zheng S, Dong K, Yu Y. Growth-promoting effect of bisphenol A on neuroblastoma *in vitro* and *in vivo*. *J Pediatr Surg* 2009;44:672–80.
- [67] Bouskine A, Nebout M, Brucker-Davis F, Benahmed M, Fenichel P. Low doses of bisphenol A promote human seminoma cell proliferation by activating PKA and PKG via a membrane G-protein-coupled estrogen receptor. *Environ Health Perspect* 2009;117:1053–8.
- [68] Dairkee SH, Seok J, Champion S, Sayeed A, Mindrinos M, Xiao W, et al. Bisphenol A induces a profile of tumor aggressiveness in high-risk cells from breast cancer patients. *Cancer Res* 2008;68:2076–80.