



Contents lists available at ScienceDirect

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Correspondence

Kisspeptin and preeclampsia



Dear Editor,

We read the recent article by Ziyaraa et al. [1] entitled “Correlation of Kisspeptin-10 level and fetal well-being in preeclamptic patients” published in the *Taiwanese Journal of Obstetrics and Gynecology* with interest. The authors reported that plasma kisspeptin-10 level was lower in preeclamptic pregnant women and inversely correlated with the severity of the disease [1]. In addition, plasma kisspeptin-10 level at either second or third trimester directly correlated with fetal birth weight in women with mild preeclampsia but inversely correlated with fetal birth weight in women with severe preeclampsia. We congratulate the success of the authors. This article is interesting and worthy discussing.

First, in the second trimester, plasma kisspeptin level was lower in women with preeclampsia and inverse correlated with the disease severity (2.30 mg/ml of controls, 2.18 mg/ml of mild preeclampsia, and 1.59 mg/ml of severe preeclampsia, respectively); however, this inverse correlation was not found in the third trimester, even though women with preeclampsia had a lower plasma level of kisspeptin consistently than women without did (2.16 mg/ml, and 2.39 mg/ml vs. 2.95 mg/ml) [1]. Plasma kisspeptin levels were dramatically elevated in women with severe preeclampsia from the second trimester to the third trimester (from 1.59 mg/ml to 2.39 mg/dl). This finding could not support the negative correlation between the plasma kisspeptin levels and severity of preeclampsia the authors made. In theory, plasma kisspeptin should be continuously low. Recently, one animal study reported that except during the pre-implantation period, kisspeptin expression was higher in the non-pregnant uterus compared to all gestational periods, indicating a pregnancy-related downregulation [2]. The other study reported that women with preeclampsia had higher placental kisspeptin expression but lower circulating serum kisspeptin levels than women without did [3], suggesting the possibility of discrepancy of kisspeptin function between local effect (paracrine and/or autocrine) and systemic effect (endocrine). If the authors could provide the data based on the longitudinal study to evaluate a correlation between plasma kisspeptin levels and gestational age, it may more easily explore the role of plasma kisspeptin. In addition, what is the cut-off value to predict the development of preeclampsia? The audience might be interested in it.

Second, fetal body weight was significantly correlated with gestational weeks [4,5]. Women with server preeclampsia have a higher risk for early delivery [6,7], indicating that their newborns might be delivered at the earlier gestational weeks, contributing to the significantly low fetal birth weight in this population. There is no exception to get the low fetal birth weight in the preeclampsia group compared to that in the normal control group. The plasma kisspeptin levels should be adjusted by gestational weeks.

In conclusion, the discovery of kisspeptin originally derived from its antimetastatic property and its role in reproductive function was a milestone in the field of reproductive biology [8]. Preeclampsia is still a biggest challenge for both physicians and pregnant women, partly because of unclear pathogenesis and partly because of a major cause of maternal, fetal and neonatal morbidity and mortality [9]. In the absence of effective treatment of preeclampsia except delivery, the identification of any new biomarker for early diagnosis of preeclampsia is welcome.

Conflicts of interest

The authors declare that there are no conflicts of interest related to the subject matter or materials discussed in this article.

Acknowledgments

This article was supported by grants from the Ministry of Science and Technology, Executive Yuan (MOST 103-2314-B-010-043-MY3), and Taipei Veterans General Hospital (V105C-096; V106C-129; V106D23-001-MY2-1; and V106A-012). We appreciate the Clinical Research Core Laboratory and the Medical Science & Technology Building of Taipei Veterans General Hospital for providing experimental space and facilities.

References

- [1] Ziyaraa MA, Hamdan FB, Mousa LR. Correlation of Kisspeptin-10 level and fetal well-being in preeclamptic patients. *Taiwan J Obstet Gynecol* 2016;55: 840–6.
- [2] Schäfer-Somi S, Ay SS, Kaya D, Sözmén M, Beceriklisoy HB, Ağaoğlu AR, et al. Kisspeptin-10 and the G protein-coupled receptor 54 are differentially expressed in the canine pregnant uterus and trophoblast cells. *Reprod Domest Anim* 2017;52(Suppl 2):123–9. <http://dx.doi.org/10.1111/rda.12818>. Epub 2016 Oct 23.
- [3] Matjila M, Millar R, van der Spuy Z, Katz A. Elevated placental expression at the maternal-fetal interface but diminished maternal circulatory kisspeptin in preeclamptic pregnancies. *Pregnancy Hypertens* 2016;6:79–87.
- [4] Tsai HJ. The birth weight of vaginal birth after cesarean section. *Taiwan J Obstet Gynecol* 2016;55:915.
- [5] Wang PH, Chen CY, Lee CN. Taiwan birth weight reference. *J Chin Med Assoc* 2014;77:553–5.
- [6] Lin LT, Wang PH, Tsui KH, Cheng JT, Cheng JS, Huang WC, et al. Increased risk of systemic lupus erythematosus in pregnancy-induced hypertension: a nationwide population-based retrospective cohort study. *Med Baltim* 2016;95: e4407.
- [7] Wang PH, Yang MJ, Chen CY, Chao HT. Endothelial cell dysfunction and preeclampsia. *J Chin Med Assoc* 2015;78:321–2.
- [8] Tsui KH, Huang BS, Wang PH. Kisspeptin system in female reproduction: a next-generation target in the manipulation of sex hormones. *J Chin Med Assoc* 2016;79:519–20.
- [9] Wang PH, Yeh CC, Chen YJ, Chen CP. Maternal serum markers and preeclampsia. *Taiwan J Obstet Gynecol* 2015;54:339–40.

Huann-Cheng Horng

Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, Taipei, Taiwan

Department of Obstetrics and Gynecology, National Yang-Ming University, Taipei, Taiwan

Institute of BioMedical Informatics, National Yang-Ming University, Taipei, Taiwan

Chang-Chin Yeh

Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, Taipei, Taiwan

Department of Obstetrics and Gynecology, National Yang-Ming University, Taipei, Taiwan

Institute of Clinical Medicine, National Yang-Ming University, Taipei, Taiwan

Peng-Hui Wang*

Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, Taipei, Taiwan

Department of Obstetrics and Gynecology, National Yang-Ming University, Taipei, Taiwan

Institute of Clinical Medicine, National Yang-Ming University, Taipei, Taiwan

Department of Medical Research, China Medical University Hospital, Taichung, Taiwan

* Corresponding author. Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, 201, Section 2 Shih-Pai Road, Taipei 112, Taiwan. Fax: +886 2 77232788.

E-mail addresses: phwang@vghtpe.gov.tw, pongpongwang@gmail.com, phwang@ym.edu.tw (P.-H. Wang).