



Review Article

Zika virus: An emerging challenge for obstetrics and gynecology



Jhe-Jhih Lin^a, Ting-Yu Chin^b, Chih-Ping Chen^{d, e, f, g, h, i}, Hong-Lin Chan^{a, **},
Tzong-Yuan Wu^{b, c, *}

^a Institute of Bioinformatics and Structural Biology, National Tsing Hua University, Hsinchu, Taiwan

^b Department of Bioscience Technology, Chung Yuan Christian University, Chungli, Taiwan

^c Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan

^d Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^e Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^f Department of Biotechnology, Asia University, Taichung, Taiwan

^g School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^h Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

ⁱ Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

ARTICLE INFO

Article history:

Accepted 20 June 2017

Keywords:

Flavivirus

Microcephaly

miRNA-34

Wnt/β-catenin signaling pathway

Zika virus

ABSTRACT

Microcephaly is a rare birth defect, however, the re-emerging mosquito and sexual transmitted *flavivirus*, Zika virus (ZIKV), had changed the situation and caused an urgent challenge for the obstetrics and gynecology. This review will brief summarize the epidemiology and virology of ZIKV. And compared the animal models that had developed for the ZIKV infections. These animal models will be benefit for the development of vaccines and anti-ZIKV drugs. Furthermore, the genes that are involved in the causation of microcephaly were also summarized. Finally, the Wnt signal is critical for the brain development as well as innate immune response. Based on previous literatures, we proposed that ZIKV-induced microcephaly might result from the influence of Wnt/β-catenin signaling pathway through the regulation of miRNA-34.

© 2017 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Microcephaly is a rare birth defect in which the baby's cranium with a significantly decrease in occipito-frontal head circumference (OFC) of greater than two standard deviations (SD) below the normative mean, whereas those at least three SD below the mean may be described as "severe microcephaly" [1]. Microcephaly is divided into primary microcephaly (apparent congenitally) and secondary microcephaly (develops postnatally) and can be genetic or acquired (caused by environmental factors) [2]. Microcephaly can cause seizures, delay in development, and impair motor function and learning abilities. According to the World Health Organization (WHO), the etiology of microcephaly is complex and occurs to be affected by several factors, including: (1) genetic

abnormalities; (2) severe malnutrition during fetal life; (3) exposure to toxic chemicals such as arsenic, mercury, and radiation; (4) infection in the womb with the TORCH (syphilis, toxoplasmosis, other infections, rubella, cytomegalovirus (CMV) and herpes simplex) group [3]. The incidence of neonatal microcephaly, as reported in birth defect registers world-wide, differs from 1.3 to 150/100,000 live births, depending upon the population type and the range of SD used to define microcephaly [1]. Although the incidence of microcephaly in general was low, however, the incidence of microcephaly has unusually risen 20-fold in Brazil since 2015 (nearly 3000 cases of microcephaly have been reported) [4]. Epidemiological studies have linked these microcephaly cases to an old, *Aedes* mosquito-transmitted *flavivirus*, Zika virus (ZIKV). This re-emerging ZIKV epidemic may change the once rare microcephaly to be a new challenge issue of obstetrics and gynecology. In this brief review, we will describe the epidemiology and virology of ZIKV and propose the possible molecular mechanism that ZIKV might "hijack" Wnt signaling pathway on the neural stem cells and specifically impair fetal brain development and lead to microcephaly.

* Corresponding author. Department of Bioscience Technology, Chung Yuan Christian University, Chungli, Taiwan. Fax: +866 3 2653599.

** Corresponding author. Fax: +886 3 5715934.

E-mail addresses: hlchan@life.nthu.edu.tw (H.-L. Chan), tywu@cycu.edu.tw (T.-Y. Wu).

The ZIKV pandemic

Before 21 century, only sporadic cases of ZIKV infection in humans were reported. However, in 2007, ZIKV emerged outside of Asia and Africa and caused the first large outbreak on Yap Island in the Federated States of Micronesia [5]. Unfortunately, a larger epidemic in French Polynesia during 2013–2014 was followed. This ZIKV outbreak in French Polynesia (FP), an overseas country of the French Republic, was estimated about 32,000 persons (11.5% of the population) with Zika-like symptoms [6,7]. Subsequently, ZIKV spread to New Caledonia [8], Japan [9], Norway [10], Easter Island [11], and continental France [12]. Recently, the ZIKV “fire” also burns to the Americas.

In 2015, there was a burst of ZIKV infection in the Americas. Brazil is the most ZIKV hit country, with estimates of 440,000–1,300,000 suspected cases of ZIKV have been reported [13,14]. Since then, ZIKV has rapidly spread far and wide across 50 other countries and territories in the Americas, including the Mexico and United States (World Map of Areas with Risk of Zika, according to the Centers for Disease Control and Prevention (CDC)). The current ZIKV epidemic provides evidence that ZIKV infection may be associated with severe neurological complications, such as Guillain–Barre syndrome (GBS) in adults in FP [15] and microcephaly in fetuses in Brazil [16]. Although, in the early 1970s, Bell et al. had showed that intracerebral infections of newborn and 5-week-old Swiss Webster mice with ZIKV can result in astrocyte hypertrophy and damage to hippocampal pyramidal neurons [17]. Furthermore, electron microscopy revealed that virions were present in both neurons and astroglial cells and morphogenesis in the endoplasmic reticulum (ER). These results have documented that ZIKV can infect and replicate in nervous system of mice, similar to other arboviruses [17].

Zika virus virology

ZIKV is an arbovirus of the Flaviviridae family, which includes dengue virus (DENV), West Nile virus (WNV), yellow fever virus, and tick-borne encephalitis virus (TBEV) [18], and is transmitted to humans by *Aedes* mosquitoes [19]. It was first isolated in Uganda in 1947 from a Rhesus monkey (strain MR-766) [20] and later the first human-infected case was detected in Nigeria in 1954 [21]. Before the 2007 outbreak in Yap, ZIKV has only been found to commonly circulate in tropical regions of Africa and Asia [22]. During outbreaks, humans serve as primary hosts for ZIKV [23], and both urban and sylvatic viral transmission have been demonstrated [24,25]. Otherwise, epizootics of ZIKV in monkeys had also been documented [26], but it is unclear whether primates are the only reservoir in the transmission of ZIKV to humans. Intriguingly, an unexpected finding is that ZIKV can be transmitted by sexual activity had been documented in 2011 [27]. And this may imply that both mosquito bites and human sexual activity could play a role in the recent ZIKV outbreak.

Like other member of the Flaviviridae family, ZIKV is constituted by a single positive sense RNA genome (+ssRNA), which is initially translated into a single polyprotein. This polyprotein is then cleaved post-translationally into individual proteins via host and viral proteases [28]. Three structural proteins named capsid (C), membrane precursor (PrM), and envelope (E) form capsids and seven nonstructural (NS) proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B and NS5) are involved in both viral transcription and replication and also modulate the host antiviral response [29,30]. Among nonstructural proteins, the glycosylated NS1 exists in two major forms, one is a membrane-bound dimer (mNS1) involved in the viral replication, and the other is a secreted hexamer (sNS1) that functions in pathogenesis and immune evasion via interacting

with immune components [31–34]. Furthermore, it can also serve as a biomarker for early detection of flavivirus infection [35]. The other six NS proteins, NS2A to NS5, are thought to be involved in the formation of a replication complex on the cytoplasmic side of the ER membrane [36]. Two major lineages of ZIKV had been revealed by phylogenetic analyses (genomic sequencing): one includes the African lineage come from the original 1947 Ugandan strain (MR-766) [37] and the Asian lineage which are confirmed to be responsible for current outbreak [38].

The common clinical manifestations of ZIKV infection resembles that of DENV and chikungunya virus (CHIKV) may include fever, headache, arthralgia, myalgia, and maculopapular rash [39]. For these complex of symptoms that also hampers clinical diagnosis. Although 80% of human ZIKV infections are asymptomatic or may not get sick at all [5]. However, cases of neurological manifestations and the GBS were reported in FP during the 2013–14 ZIKV outbreak [12]. GBS is a rare autoimmune disorder in which the body's immune system attacks the peripheral nerves [40]. Often, the first symptoms of this disorder are weakness and tingling sensations in legs [41]. The exact cause of GBS is unknown and a new paradox was added that why this neurological disorder can be caused by ZIKV infection. However, it is learned that GBS is frequently preceded by an infectious illness such as acute respiratory or gastrointestinal infection [42]. There were 42 cases of GBS occurring during ZIKV outbreak in FP was recorded [43]. The observation of GBS in ZIKV cases implied an increase in the potential clinical severity of the disease [44]. More bizarre is that the Brazil Health Ministry reported a dramatically increased number of neonatal microcephaly cases (approximately 20 times higher than previous years) in the northeast region of Brazil and concerned this unusual epidemic linked with ZIKV [16]. And after Brazil raised the alarm, health officials in FP also identified more than a dozen babies born with neural defects [4]. Based on these epidemic observations, an emerging issue to clarify is that: Does the ZIKV is the only pathogen that lead to neurological manifestations, especially the microcephaly?

Zika virus caused microcephaly in animals and human

As aforementioned, the unusual increased microcephaly cases in the “hot zone” of ZIKV in Brazil, which may imply that ZIKV is the pathogenic factor. Moreover, causality relationship between the ZIKV epidemic and microcephaly had been experimentally confirmed and mentioned in a clinical case.

A study was conducted to examine the effects of ZIKV infection in neurospheres and brain organoids that generated from human iPSC-derived neural stem cells. This *in vitro* model demonstrates that ZIKV directly infects the brain cells and impairs their viability and grows in human neurospheres and brain organoids. These results implied that ZIKV interferes neurogenesis during human brain development [45]. To further confirm the observations of ZIKV infection in an *in vitro* brain organoids model, the establishment of animal models is necessary and critical. And these animal models should also be benefit for the development of the anti-ZIKV drugs and vaccines. At present, two kind of murine models of ZIKV infection have been established and published, including adult models [46,47] and pregnancy models [48–50], are summarized in Table 1. Moreover, several non-human primate models to study ZIKV are also under way [51]. These studies provide solid evidences that ZIKV can replicate in the embryonic brain and cause severe fetal abnormalities, which is consistent with the reported Zika-related microcephaly cases during the 2015–2016 American outbreak [39].

To test whether ZIKV can infect the embryonic mouse brain, ZIKV^{SZ01} (Asian ZIKV strain) were directly injected into one side of

Table 1

The murine models used in clarify linkage between ZIKV and microcephaly.

ZIKV Strain used	Strain of mice	Inoculation routes	Virus titers	Infected cells (or tissue) identified	Reference
Adult model					
1. Asian-lineage ZIKV (FSS13025)	3-week-old, 5-week-old, or 11-week-old A129 mice 3-week-old AG129	Intraperitoneal (IP) route Intradermal (ID) route	1.0×10^4 PFU/mouse 1.0×10^5 PFU/mouse	Heart, lung, liver, kidney, and muscle Spleen, brain, and testes	[47]
2. Asian-lineage ZIKV (H/PF/2013) African-lineage ZIKV (MR-766, Uganda)	4- to 6-week-old <i>Ifnar1</i> ^{-/-} mice <i>Irf3</i> ^{-/-} <i>Irf5</i> ^{-/-} <i>Irf7</i> ^{-/-} TKO mice (C57BL/6 background)	Subcutaneous (footpad) route Intravenous (IV) route	1.0×10^2 FFU/mouse 1.0×10^3 FFU/mouse 1.0×10^4 FFU/mouse	Serum, spleen, liver, and kidney Brain, spinal cord, and testes	[46]
Pregnant model					
1. Asian-lineage ZIKV (SZ01/2016/China)	E13.5 ICR mice	Directly injected into one side of the cerebroventricular space/LV	6.5×10^2 PFU/mouse	Fetal brain Neural progenitor cells (NPCs)	[49]
2. Asian-lineage ZIKV (H/PF/2013)	E6.5–7.5 <i>Ifnar1</i> ^{+/-} mice (C57BL/6 background)	Subcutaneous (footpad) route	1.0×10^3 FFU/mouse	Serum, spleen, fetal brain, and placenta Trophoblast cells (including glycogen trophoblasts and spongiotrophoblasts)	[50]
3. Asian-lineage ZIKV (Brazilian ZIKV strain)	Day 10–13 of gestation SJL mice	Intravenous (IV) route	2.0×10^2 PFU/mouse 8.0×10^9 PFU/mouse 2.0×10^{11} PFU/mouse	Brain Cortical progenitor cells	[48]

the cerebroventricular space/lateral ventricle (LV) of embryonic day 13.5 (E13.5) ICR mouse brains and inspected 3–5 days later [49]. Real-time PCR and immunohistochemistry verified that the brains were readily infected and the number of ZIKV RNA copies increased in 300 times was detected 3 days after infection. Besides, the sizes of ZIKV infected embryonic brains are smaller as compared to those of their mock infected control. Similar to the *in vitro* human brain organoids model, this *in vivo* model also found that ZIKV leads to cell-cycle arrest, apoptosis, and inhibition of neural progenitor cell (NPC) differentiation in the infected embryonic brain. These results may provide pathological evidences that ZIKV infection will cause cortical thinning and microcephaly [49].

During the period of 2013–2015 Brazil outbreak, ZIKV has been linked with microcephaly and other severe neurological diseases, such as GBS [52,53]. Besides the suspected clinical evidences, direct experimental data confirming that the ZIKV^{BR} (Brazilian ZIKV strain which was isolated from a baby born with microcephaly in Paraiba, a state in the Northeast Region of Brazil) is the only pathogen that causes microcephaly and other neurological syndromes was also reported recently [48]. To understand the relationship between ZIKV^{BR} infection and birth defects, a pregnant mouse model (10–13 days of gestation) in which SJL (a immunocompetent mice with high susceptibility to several different viral infections [54]) and C57BL/6 (control) mice (6–8 weeks of age) were infected with ZIKV^{BR} via an intravenous route [48]. These results clearly demonstrated that ZIKV^{BR} can cross the placenta and infect fetuses and display intrauterine growth restriction (IUGR), which caused congenital malformation (including microcephaly) in the pregnant SJL mice. Additionally, ZIKV^{BR} selectively infects cortical progenitor cells of fetal brain, inducing cell death through apoptosis and autophagy as well as impairing neurodevelopment [48].

Apart from aforementioned studies, two mouse models, one with genetic knockout approach and another with antibody mediated “functional knock-down”, were also developed to study in utero ZIKV transmission and infection [50]. In the gene knockout approach, pregnant females *IFNαβR*^{-/-} mice that lacking type I interferon (IFNs) signaling were mated to wild-type (WT) males (C57BL/6) to generate *IFNαβR*^{+/-} (heterozygous) pups which display similar immune status to human fetuses, whereas in a complementary approach using pregnant WT females treated with an anti-ifnar-blocking antibody (MAR1-5A3) to functional knock-down of IFN-α receptor 1 by intraperitoneal injection. Using these mice as an animal model for ZIKV infection, pregnant dams were inoculated with ZIKV (H/PF/2013), which was isolated from a

clinical patient in FP, on E6.5 or E7.5 via a subcutaneous (footpad) route. These results showed that ZIKV can replicate in placental trophoblasts and induce cellular apoptosis, which is involved in disruption of the placental barrier, as well as subsequent trans-placental infection of the developing fetus. Thus, most fetuses suffered in utero demise, while surviving fetuses exhibited significant IUGR and high levels of viral RNA in fetal brains [50].

Similarly, there are two recent reports using IFN-related transgenic mice models to study the association between ZIKV infection and neurological disorders [46,47]. To characterize ZIKV in immunocompetent mice, Lazarar et al. subcutaneously injected with 10^3 or 10^4 FFU of ZIKV strain H/PF/2013 into 4- to 6-week-old WT C57BL/6 mice and transgenic mice lacking key components of innate antiviral immunity. *Ifnar1*^{-/-} mice which lacking the interferon receptor cannot respond to IFN-α/β and *Irf3*^{-/-} *Irf5*^{-/-} *Irf7*^{-/-} triple knockout (TKO) mice which produce almost no IFN-α/β were highly susceptible to ZIKV infection and presented with neurological disease [46]. In contrast, single *Irf3*^{-/-}, *Irf5*^{-/-}, and *Mavs*^{-/-} knockout mice exhibited no signs of illness when infected with ZIKV. Noteworthy, *Ifnar1*^{-/-} mice sustained high viral loads in the brain and spinal cord and developed highest levels of ZIKV infection in the testes, which is relevant to sexual transmission of ZIKV. These results provide solid evidence implying that ZIKV infection can affect neurodevelopment in human fetuses and can be contracted either by mosquito biting or through sexual transmission [46]. Additional, immunocompromised mice A129 (lacking the type I IFN-receptor) and AG129 (lacking both type I and type II IFN-receptors) have also observed similar results, when challenged with a low-passage Cambodian isolate (FSS13025), an Asian ZIKV strain [47]. These animal models, as summarized in Table 1, will be valuable for the study of ZIKV pathogenesis as well as assessing the efficacy of therapeutic drugs and vaccines.

Besides the animal studies implied the association between ZIKV infection and microcephaly, a direct human case of ZIKV in Brazil was recorded in 2016 [39]. In this case, an expectant mom presented a febrile illness with rash was probably infected with ZIKV at the end of the first trimester of pregnancy and the fetus exhibited microcephaly and severe brain injury. This report provides the evidence that fetal brain abnormalities related to ZIKV infection with vertical transmission [39]. Ultrasonographic examination that was performed at 32 weeks of gestation showed small head circumference (microcephaly) and numerous calcifications in the brain and placenta. A fetal autopsy was conducted 3 days after the end of the pregnancy which showed several brain

abnormalities including micrencephaly (an abnormally small brain), internal hydrocephalus, almost complete agyria, spotty calcifications in the cortex and subcortical white matter, cortical displacement, and focal inflammation. In addition, reverse transcriptase-polymerase chain reaction (RT-PCR) assays showed high levels of ZIKV in the fetal brain tissue. Furthermore, the complete genome of this ZIKV was recovered from the fetal brain, named ZIKV strain Bahia, Brazil (KU527068) is consistent with the strain circulating in Brazil comes from the Asian lineage [39]. This study also showed that placental calcification and low placental–fetal weight ratio of 0.136 (<3rd percentile), consistent with previous studies that ZIKV could damage the placenta [48,50]. More interesting, there were no virus and no pathological effects in other fetal organs, suggests a specific neurotropism of the ZIKV [39]. This open a question that the neurotropism of the virus is restricted through the entry pathways (e.g., with a neuronal specific receptor) or the specific replication strategy of the virus in the brain that may be interfered with the production of ZIKV in other tissues.

Genes that correlates microcephaly

Before the 2015 outbreak of ZIKV in Brazil which is indeed causing the severe epidemic of microcephaly, this abnormal brain development is an unusual event in obstetrics and gynecology [15]. However, there are many genetic studies that identified the genes that may associated with microcephaly and are summarized in Table 2. Thus, it is interesting to learn whether the ZIKV infection have any effect on the expression of these microcephaly associated genes. In the “sequence everything” approach of next generation sequencing (NGS) era, it is easy and straightforward to search for genes that were altered upon ZIKV infection [55–57]. Global gene expression analysis of infected brain organoids might provide insights into the link between ZIKV infection and microcephaly and further to figure out the potential underlying mechanisms and management of ZIKV-related pathological effects during brain development [58,59]. Li et al. (2016) carried out global transcriptome analyses determined by RNA-seq and identified several differentially expressed genes at 3 days in ZIKV-infected ICR mouse [49]. Gene ontology (GO) analyses showed that genes associated with microcephaly, including ASPM, CASC5, MCPH1, RBBP8, STIL, and TBR2 (all are listed in Table 2) were significantly downregulated. In addition, genes involved in organ development as well as participated in cell proliferation, differentiation, and migration were also significantly downregulated. In contrast, GO analysis revealed enrichment of upregulated genes related to immune response and apoptosis pathways. Among these upregulated genes, several genes involved in production of cytokines and the response to cytokines were notable. Thus, it is suggested that cytokines may play a significant role in the pathogenesis of ZIKV infection [49].

Zika virus regulates the Wnt signal during brain development-a hypothesis

The Wnt/ β -catenin signaling pathway has been shown to be involved in regulation of development, cellular growth, cell–cell interactions, tumorigenesis, and stem cell biology [60–64]. Wnt signaling is initiated through the binding of soluble Wnt ligands to cell surface receptors of the Frizzled (Fz) family and co-receptor of the low-density lipoprotein receptor-related protein (LRP) family. This receptor-ligand complex recruits the scaffold protein, Dishevelled (DVL), and in turn recruits the Axin-glycogen synthase kinase 3 β (GSK3 β) complex, resulting in the phosphorylation of LRP by GSK3 β [65–67]. This signalosome-mediated recruitment of more Axin from the β -catenin destruction complex to the plasma membrane [68,69]. The “ β -catenin destruction complex” is a multi-

domain scaffold phospho-protein complex, that includes Axin, adenomatous polyposis coli (APC), glycogen synthase kinase 3 β (GSK3 β), Casein Kinase 1 (CK1), protein phosphatase 2A (PP2A), and E3-ubiquitin ligase [70]. Recruitment of Axin lead to disassembly of the destruction complex resulting in the repression of GSK3 β activity. Inhibition of GSK3 β -mediated phosphorylation leads to stabilization and accumulation of cytosolic β -catenin [71]. The rise of cytoplasmic levels of β -catenin then translocates to the nucleus where interacting with the TCF/LEF transcription factors and thereby switch on Wnt-target gene transcription. In the absence of Wnt, GSK3 β constitutively phosphorylates β -catenin, marking it for ubiquitination and subsequent proteasome-mediated degradation [72,73].

Previous studies have explored the role of Wnt signaling in the regulation of cerebral cortical size [74,75]. To examine whether Wnt signal could regulate mammalian brain development, a genetic approach is employed to generate transgenic mice that overexpress a stabilized form of β -catenin in neuronal progenitor cells [74]. β -catenin can act as an important component of adherens junctions [76], also serve as a transcriptional co-activator of Wnt target gene expression [77], and is highly expressed in mammalian neural progenitors [74]. These studies showed that over activating β -catenin influences cell fate decisions during brain development and leads the neural precursors to re-enter the cell cycle after mitosis. Thus, overexpression of β -catenin results in enlarged brain size which has a greatly increased cerebral cortical surface area without increased cortical thickness [74]. In contrast, inactivation of Wnt signals leads to specific developmental brain defects, including reduction of hippocampal development [62] or absence of midbrain and cerebellum [60]. Based on these observations, we hypothesized that ZIKV-induced microcephaly might result from the influence of Wnt signaling pathway.

Several studies have found that microRNAs (miRNAs) might regulate components of the Wnt signaling pathways to affect various cellular processes during embryo development [78]. miRNAs are small (20- to 23-nucleotide [nt]) and noncoding RNAs that play significant roles in regulating a broad of cellular processes through the posttranscriptional modulation of gene expression. More interestingly, there are reports indicating that viral infection induces significant changes in the expression levels of cellular miRNAs [79–83]. These observations may imply that the manipulation Wnt signaling pathways by the virus or the host response to viral infection through miRNAs. Recently, a microRNA screen approach was designed to understand the relationship between flavivirus infection and Wnt signaling pathway [84]. These results show that miR-34 family have the ability to repress Wnt signaling and exhibit antiviral effects in transfected cells infected with flaviviruses, including DENV, WNV, and ZIKV (strain PRVABC059). Further studies demonstrate that miR-34a repression the Wnt signaling, which inhibits the phosphorylation of GSK3 β , results in a direct interaction between GSK3 β and TANK-binding protein 1 (TBK1) that promotes interferon regulatory factor 3 (IRF3) phosphorylation/homodimerization and subsequently allows its translocation into the nucleus to activate the interferon responsive genes as well as to promote antiviral responses [84]. Consistent with previous established concepts that the cross talk between the Wnt pathway and the interferon-based innate immune pathway (the type I IFN signaling pathway) during viral infection [85–89]. Using locked nucleic acid (LNA)-anti-miR to knock-down endogenous miR-34a a significant increase in viral replication was observed, suggesting that endogenous miR-34a might function to suppress flaviviral replication through its ability to dampen Wnt signaling, allowing GSK3 β -TBK1-mediated activation of IRF3 in response to pathogen-associated molecular pattern (PAMP) detection. However, the endogenous levels of miR-34a was unaffected by DENV

Table 2

The genes that are correlated to microcephaly.

Gene	Protein	Function	Chromosome location	References
<i>MCPH1</i> (<i>BRIT1</i>)	Microcephalin	DNA damage repair; chromosome condensation;	8p23	[92–94]
<i>WDR62</i> (<i>MCPH2</i>)	WDR62	Transcriptional regulation of DNA damage genes Play a role in cerebral cortical development; For centriole duplication	19q13.12	[95,96]
<i>CDK5RAP2</i> (<i>MCPH3</i> , <i>Cep215</i>)	CDK5RAP2	Regulating microtubule dynamics; Centrosome maturation and cohesion	9q33.2	[97–99]
<i>KNL1</i> (<i>CASC5</i> , <i>MCPH4</i>)	KNL1 (<i>CASC5</i>)	Essential for spindle-assembly checkpoint signaling; For correct chromosome alignment; involved in centriol duplication	15q15.1	[100–102]
<i>ASPM</i> (<i>ASP</i> , <i>MCPH5</i>)	ASPM	Orientation and organization of mitotic spindles During embryonic neurogenesis	1q31.3	[103–105]
<i>CENPJ</i> (<i>CPAP</i> , <i>MCPH6</i>)	CENPJ	Centriole biogenesis and length control; Microtubule dynamics	13q12.12	[106–108]
<i>STIL</i> (<i>SIL</i> , <i>MCPH7</i>)	STIL	Spindle organization; Hh/Shh signaling; Cell cycle progression	1p33	[109–111]
<i>CEP152</i> (<i>MCPH9</i>)	CEP152	Necessary for centrosome duplication	15q21.1	[112]
<i>CENPE</i> (<i>MCPH13</i>)	CENP-E	For the maintenance of chromosomal stability; For the movement of chromosomes toward the metaphase	4q24-q25	[113]
<i>SASS6</i> (<i>SAS6</i> , <i>MCPH14</i>)	SASS6	For centrosome biogenesis and duplication	1p21.2	[114,115]
<i>MFSD2A</i> (<i>NLS1</i> , <i>MCPH15</i>)	NLS1	For blood–brain barrier (BBB) formation and function	1p34.2	[116,117]
<i>ANKLE2</i> (<i>MCPH16</i>)	ANKLE2	Mitotic nuclear envelope reassembly; Facilitating nuclear envelope assembly	12q24.33	[118,119]
<i>CEP63</i> (<i>SCKL6</i>)	CEP63	For normal spindle assembly; plays a key role in mother-centriole- Dependent centriole duplication and in DNA damage response	3q22.2	[120,121]
<i>DYRK1A</i> (<i>DYRK</i>)	DYRK1A	Regulating cell proliferation; involved in brain development	21q22.13	[122,123]
<i>EOMES</i> (<i>TBR2</i>)	eomesodermin (<i>TBR2</i>)	Required for trophoblast development and mesoderm formation; Key regulator of neurogenesis in the SVZ	3p24.1	[124–126]
<i>KIF11</i> (<i>EG5</i>)	KIF11	Establishing a bipolar spindle during mitosis; Chromosome positioning; centrosome separation	10q23.33	[127,128]
<i>RBBP8</i> (<i>CTIP</i>)	RBBP8	Bind directly to retinoblastoma protein, regulates cell proliferation; DNA damage signaling and repair	18q11.2	[129–131]
<i>WDFY3</i> (<i>ALFY</i>)	WDFY3	For selective autophagy (aggrephagy); Formation and degradation of cytoplasmic ubiquitin-positive inclusions	4q21.23	[132,133]

Abbreviations. MCPH: Autosomal recessive primary microcephaly; MCPH1: Microcephalin; BRIT1: BRCA1 C terminus-repeat inhibitor of human telomerase expression 1; WDR62: WD repeat domain 62; CDK5RAP2: CDK5 regulatory subunit associated protein 2; KNL1: kinetochore scaffold 1; CASC5: Cancer susceptibility candidate 5; ASPM: abnormal spindle microtubule assembly; CENPJ: Centromere protein J; CPAP: Centrosomal P4.1-associated protein; STIL: SCL/TAL1 interrupting locus; CEP152: Centrosomal protein 152 (kDa); CENPE: centromere protein E; SASS6: SAS-6 centriolar assembly protein; MFSD2A: major facilitator superfamily domain containing 2A; NLS1: Sodium-dependent LPC symporter 1; ANKLE2: ankyrin repeat and LEM domain containing 2; WDFY3: WD repeat and FYVE domain containing 3; ALFY: Autophagy-Linked FYVE Protein; CEP63: centrosomal protein 63; SCKL6: Seckel syndrome-6; DYRK1A: dual specificity tyrosine phosphorylation regulated kinase 1A; KIF11: Kinesin family member 11; RBBP8: Retinoblastoma-binding protein 8; CTIP: CtBP-interacting protein; EOMES: Eomesodermin; TBR2: T-box brain protein 2.

infection [84]. It is interesting to know whether the ZIKV infection can up-regulate the levels of miR-34a to down regulation Wnt signaling pathway. Based on these observations mentioned above, we hypothesized that ZIKV infection might contribute to microcephaly through (1) induction of changes in miRNA expression levels, like up-regulate the levels of miR-34a. (2) This changes might repression of Wnt/ β -catenin signaling pathway. (3) Inactivation of Wnt signals would result in brain defects in the fetus during pregnancy. According this model, inactivation of Wnt signals would up-regulation of IFN-related genes (e.g. type I IFNs and interferon-stimulated genes (ISGs)) will promote antiviral activity for women in pregnancy, but will cause brain developmental defect, such as the microcephaly, in infants.

However, other mechanisms may also exist to explain the causal relationship between the ZIKV infection and microcephaly. For example, computational prediction of miRNAs (genome-wide) binding sites in the prototypic ZIKV (MR-766) whole genome had discovered six miRNAs, miR-627–5, miR-4646, miR-1304, miR-6771, miR-4528, and miR-3198 that share mutual homology with the microcephaly related genetic sequences MCPH1, 2, 3, 4, 5, 6, 7, and 9 (see Table 2), respectively. These findings also suggest that ZIKV may modulate host miRNAs that are involved in regulating MCPH genes during fetal brain development, resulting in microcephaly [90]. Noteworthy, miR-1304, which has homology to six

different MCPH genes (MCPH 3, 4, 8, 9, 10, and 12), might be associated with gene disorders, neurological diseases, and nervous system development had been documented [91].

Conclusion

The re-emerging mosquito and sexual transmitted ZIKV is an urgent challenge for the obstetrics and gynecology. ZIKV and the so-called STORCH factors will be the main reasons that compromise brain development in utero and cause microcephaly. Thanks to the modern biotechnology, various animal models for the ZIKV infection have been established and will facilitate the development of vaccine and anti-ZIKV drugs. We proposed that ZIKV-induced microcephaly might result from the influence of Wnt signaling pathway through the regulation of miRNA and this may be verified through experimental test and also provided a route for the antiviral research. A critical issue to be answered will be: If a woman once has contracted with ZIKV but recovered, are future pregnancies safe?

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

Acknowledgments

This work was supported by grants (MOST- 106-2321-B-033-001 -) to T. Y. Wu from the Ministry of Science and Technology of Taiwan.

References

- [1] Kaindl AM, Passemard S, Kumar P, Kraemer N, Issa L, Zwirner A, et al. Many roads lead to primary autosomal recessive microcephaly. *Prog Neurobiol* 2010;90:363–83.
- [2] Woods CG, Parker A. Investigating microcephaly. *Arch Dis Child* 2013;98:707–13.
- [3] Alvarado-Socarras JL, Rodriguez-Morales AJ. Etiological agents of microcephaly: implications for diagnosis during the current Zika virus epidemic. *Ultrasound Obstet Gynecol* 2016;47:525–6.
- [4] Vogel G. EMERGING DISEASES. A race to explain Brazil's spike in birth defects. *Science* 2016;351:110–1.
- [5] Duffy MR, Chen T-H, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med* 2009;360:2536–43.
- [6] Cao-Lormeau V-M, Roche C, Teissier A, Robin E, Berry A-L, Mallet H-P, et al. Zika virus, French polynesia, South pacific, 2013. *Emerg Infect Dis* 2014;20:1085–6.
- [7] Musso D, Gubler DJ. Zika virus. *Clin Microbiol Rev* 2016;29:487–524.
- [8] Dupont-Rouzeyrol M, O'Connor O, Calvez E, Daurès M, John M, Grangeon J-P, et al. Co-infection with Zika and Dengue viruses in 2 patients, New Caledonia, 2014. *Emerg Infect Dis* 2015;21:381–2.
- [9] Kutsuna S, Kato Y, Takasaki T, Moi M, Kotaki A, Uemura H, et al. Two cases of Zika fever imported from French Polynesia to Japan, December 2013 to January 2014 [corrected]. *Euro Surveill* 2014;19.
- [10] Wæhre T, Maagard A, Tappe D, Cadar D, Schmidt-Chanasit J. Zika virus infection after Travel to Tahiti, December 2013. *Emerg Infect Dis* 2014;20:1412–4.
- [11] Tognarelli J, Ulloa S, Villagra E, Lagos J, Aguayo C, Fasca R, et al. A report on the outbreak of Zika virus on Easter Island, South Pacific, 2014. *Arch Virol* 2016;161:665–8.
- [12] Ios S, Mallet H-P, Leparç Goffart I, Gauthier V, Cardoso T, Herida M. Current Zika virus epidemiology and recent epidemics. *Med Mal Infect* 2014;44:302–7.
- [13] Zanluca C, Melo VCA de, Mosimann ALP, Santos GIVD, Santos CNDD, Luz K. First report of autochthonous transmission of Zika virus in Brazil. *Mem Inst Oswaldo Cruz* 2015;110:569–72.
- [14] Hennessey M. Zika virus spreads to new areas — region of the Americas, May 2015–January 2016. *MMWR Morb Mortal Wkly Rep* 2016;65.
- [15] Chan JFW, Choi GKY, Yip CCY, Cheng VCC, Yuen K-Y. Zika fever and congenital Zika syndrome: an unexpected emerging arboviral disease. *J Infect* 2016;72:507–24.
- [16] Schuler-Faccini L. Possible association between Zika virus infection and microcephaly — Brazil, 2015. *MMWR Morb Mortal Wkly Rep* 2016;65.
- [17] Bell TM, Field EJ, Narang HK. Zika virus infection of the central nervous system of mice. *Arch Für Gesamte Virusforsch* 1971;35:183–93.
- [18] Gould EA, Solomon T. Pathogenic flaviviruses. *Lancet* 2008;371:500–9.
- [19] Digne CT, Diallo D, Faye O, Ba Y, Faye O, Gaye A, et al. Potential of selected Senegalese *Aedes* spp. mosquitoes (Diptera: Culicidae) to transmit Zika virus. *BMC Infect Dis* 2015;15:492.
- [20] Dick GWA, Kitchen SF, Haddow AJ. Zika virus. I. Isolations and serological specificity. *Trans R Soc Trop Med Hyg* 1952;46:509–20.
- [21] Macnamara FN. Zika virus: a report on three cases of human infection during an epidemic of jaundice in Nigeria. *Trans R Soc Trop Med Hyg* 1954;48:139–45.
- [22] Faye O, Freire CCM, Iamarino A, Faye O, de Oliveira JVC, Diallo M, et al. Molecular evolution of Zika virus during its emergence in the 20(th) century. *PLoS Negl Trop Dis* 2014;8:e2636.
- [23] Staples JE, Dziuban EJ, Fischer M, Cragan JD, Rasmussen SA, Cannon MJ, et al. Interim guidelines for the evaluation and testing of infants with possible congenital Zika virus infection — United States, 2016. *MMWR Morb Mortal Wkly Rep* 2016;65:63–7.
- [24] Grard G, Caron M, Mombo IM, Nkoghe D, Mbouli Ondo S, Jiolle D, et al. Zika virus in Gabon (Central Africa)—2007: a new threat from *Aedes albopictus*? *PLoS Negl Trop Dis* 2014;8:e2681.
- [25] Berthet N, Nakouné E, Kamgang B, Selekon B, Descorps-Declère S, Gessain A, et al. Molecular characterization of three Zika flaviviruses obtained from sylvatic mosquitoes in the Central African Republic. *Vector Borne Zoonotic Dis* 2014;14:862–5.
- [26] Foy BD, Kobylinski KC, Foy JLC, Blitvich BJ, Travassos da Rosa A, Haddow AD, et al. Probable non-vector-borne transmission of Zika virus, Colorado, USA. *Emerg Infect Dis* 2011;17:880–2.
- [27] Foy BD, Kobylinski KC, Foy JLC, Blitvich BJ, Travassos da Rosa A, Haddow AD, et al. Probable non-vector-borne transmission of Zika virus, Colorado, USA. *Emerg Infect Dis* 2011;17:880–2.
- [28] Murray CL, Jones CT, Rice CM. Architects of assembly: roles of Flaviviridae non-structural proteins in virion morphogenesis. *Nat Rev Microbiol* 2008;6:699–708.
- [29] Lindenbach BD, Thiel HJ, Rice CM, Lindenbach BD, Thiel H, Rice C, et al. Flaviviridae: the viruses and their replication. 2007.
- [30] Cunha MS, Esposito DLA, Rocco IM, Maeda AY, Vasami FGS, Nogueira JS, et al. First complete genome sequence of Zika virus (Flaviviridae, Flavivirus) from an autochthonous transmission in Brazil. *Genome Announc* 2016;4.
- [31] Young PR, Hilditch PA, Bletchly C, Halloran W. An antigen capture enzyme-linked immunosorbent assay reveals high levels of the dengue virus protein NS1 in the sera of infected patients. *J Clin Microbiol* 2000;38:1053–7.
- [32] Chung KM, Liszewski MK, Nybakken G, Davis AE, Townsend RR, Fremont DH, et al. West Nile virus nonstructural protein NS1 inhibits complement activation by binding the regulatory protein factor H. *Proc Natl Acad Sci U S A* 2006;103:19111–6.
- [33] Suthar MS, Diamond MS, Gale M. West Nile virus infection and immunity. *Nat Rev Microbiol* 2013;11:115–28.
- [34] Youn S, Ambrose RL, Mackenzie JM, Diamond MS. Non-structural protein-1 is required for West Nile virus replication complex formation and viral RNA synthesis. *Virol J* 2013;10:339.
- [35] Cecchetto J, Fernandes FCB, Lopes R, Bueno PR. The capacitive sensing of NS1 Flavivirus biomarker. *Biosens Bioelectron* 2017;87:949–56.
- [36] Song H, Qi J, Haywood J, Shi Y, Gao GF. Zika virus NS1 structure reveals diversity of electrostatic surfaces among flaviviruses. *Nat Struct Mol Biol* 2016;23:456–8.
- [37] Haddow AD, Schuh AJ, Yasuda CY, Kasper MR, Heang V, Huy R, et al. Genetic characterization of Zika virus strains: geographic expansion of the Asian lineage. *PLoS Negl Trop Dis* 2012;6:e1477.
- [38] Wikan N, Smith DR. Zika virus: history of a newly emerging arbovirus. *Lancet Infect Dis* 2016;16:e119–126.
- [39] Malakar J, Korva M, Tul N, Popović M, Poljšak-Prijatelj M, Mraz J, et al. Zika virus associated with microcephaly. *N Engl J Med* 2016;374:951–8.
- [40] Oehler E, Watrin L, Larre P, Leparç-Goffart I, Lasteré S, Valour F, et al. Zika virus infection complicated by Guillain-Barré syndrome — case report, French Polynesia, December 2013. *Euro Surveill* 2014;19.
- [41] Wijedicks EFM, Klein CJ. Guillain-Barré syndrome. *Mayo Clin Proc* 2017;92:467–79.
- [42] Vucic S, Kiernan MC, Cornblath DR. Guillain-Barré syndrome: an update. *J Clin Neurosci* 2009;16:733–41.
- [43] Watrin L, Ghawché F, Larre P, Neau J-P, Mathis S, Fournier E. Guillain-Barré syndrome (42 cases) occurring during a Zika virus outbreak in French Polynesia. *Medicine (Baltimore)* 2016;95.
- [44] Roth A, Mercier A, Lepers C, Hoy D, Duituturaga S, Benyon E, et al. Concurrent outbreaks of dengue, chikungunya and Zika virus infections — an unprecedented epidemic wave of mosquito-borne viruses in the Pacific 2012–2014. *Euro Surveill* 2014;19.
- [45] Garcez PP, Loliola EC, Costa RM da, Higa LM, Trindade P, Delvecchio R, et al. Zika virus impairs growth in human neurospheres and brain organoids. *Science* 2016;352:816–8.
- [46] Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, et al. A mouse model of Zika virus pathogenesis. *Cell Host Microbe* 2016;19:720–30.
- [47] Rossi SL, Tesh RB, Azar SR, Muruato AE, Hanley KA, Auguste AJ, et al. Characterization of a novel murine model to study Zika virus. *Am J Trop Med Hyg* 2016;94:1362–9.
- [48] Cugola FR, Fernandes IR, Russo FB, Freitas BC, Dias JLM, Guimarães KP, et al. The Brazilian Zika virus strain causes birth defects in experimental models. *Nature* 2016;534:267–71.
- [49] Li C, Xu D, Ye Q, Hong S, Jiang Y, Liu X, et al. Zika virus disrupts neural progenitor development and leads to microcephaly in mice. *Cell Stem Cell* 2016;19:120–6.
- [50] Miner JJ, Cao B, Govero J, Smith AM, Fernandez E, Cabrera OH, et al. Zika virus infection during pregnancy in mice causes placental damage and fetal demise. *Cell* 2016;165:1081–91.
- [51] Rossi SL, Vasilakis N. Modeling Zika virus infection in mice. *Cell Stem Cell* 2016;19:4–6.
- [52] Aragao M de FV, Linden V van der, Brainer-Lima AM, Coeli RR, Rocha MA, Silva PS da, et al. Clinical features and neuroimaging (CT and MRI) findings in presumed Zika virus related congenital infection and microcephaly: retrospective case series study. *BMJ* 2016;353:i1901.
- [53] Cao-Lormeau V-M, Blake A, Mons S, Lastère S, Roche C, Vanhomwegen J, et al. Guillain-Barré Syndrome outbreak associated with Zika virus infection in French Polynesia: a case-control study. *Lancet* 2016;387:1531–9.
- [54] Spindler KR, Fang L, Moore ML, Hirsch GN, Brown CC, Kajon A. SJL/J mice are highly susceptible to infection by mouse adenovirus type 1. *J Virol* 2001;75:12039–46.
- [55] Calvet G, Aguiar RS, Melo ASO, Sampaio SA, Filippis I de, Fabri A, et al. Detection and sequencing of Zika virus from amniotic fluid of fetuses with microcephaly in Brazil: a case study. *Lancet Infect Dis* 2016;16:653–60.
- [56] Liu L, Wu W, Zhao X, Xiong Y, Zhang S, Liu X, et al. Complete genome sequence of Zika virus from the first imported case in mainland China. *Genome Announc* 2016;4.
- [57] Sardi SI, Somasekar S, Naccache SN, Bandeira AC, Tauro LB, Campos GS, et al. Coinfections of Zika and Chikungunya viruses in Bahia, Brazil, identified by metagenomic next-generation sequencing. *J Clin Microbiol* 2016;54:2348–53.

- [58] Tang H, Hammack C, Ogden SC, Wen Z, Qian X, Li Y, et al. Zika virus infects human cortical neural progenitors and attenuates their growth. *Cell Stem Cell* 2016;18:587–90.
- [59] Zhang F, Hammack C, Ogden SC, Cheng Y, Lee EM, Wen Z, et al. Molecular signatures associated with ZIKV exposure in human cortical neural progenitors. *Nucleic Acids Res* 2016;44:8610–20.
- [60] McMahon AP, Bradley A. The Wnt-1 (int-1) proto-oncogene is required for development of a large region of the mouse brain. *Cell* 1990;62:1073–85.
- [61] Moon RT, Brown JD, Torres M. WNTs modulate cell fate and behavior during vertebrate development. *Trends Genet* 1997;13:157–62.
- [62] Lee SM, Tole S, Grove E, McMahon AP. A local Wnt-3a signal is required for development of the mammalian hippocampus. *Development* 2000;127:457–67.
- [63] Brault V, Moore R, Kutsch S, Ishibashi M, Rowitch DH, McMahon AP, et al. Inactivation of the beta-catenin gene by Wnt1-Cre-mediated deletion results in dramatic brain malformation and failure of craniofacial development. *Dev Camb Engl* 2001;128:1253–64.
- [64] Van Camp JK, Beckers S, Zegers D, Van Hul W. Wnt signaling and the control of human stem cell fate. *Stem Cell Rev* 2014;10:207–29.
- [65] Wong H-C, Bourdelas A, Krauss A, Lee H-J, Shao Y, Wu D, et al. Direct binding of the PDZ domain of Dishevelled to a conserved internal sequence in the C-terminal region of Frizzled. *Mol Cell* 2003;12:1251–60.
- [66] Bilic J, Huang Y-L, Davidson G, Zimmermann T, Cruciat C-M, Bienz M, et al. Wnt induces LRP6 signalosomes and promotes dishevelled-dependent LRP6 phosphorylation. *Science* 2007;316:1619–22.
- [67] Zeng X, Huang H, Tamai K, Zhang X, Harada Y, Yokota C, et al. Initiation of Wnt signaling: control of Wnt coreceptor Lrp6 phosphorylation/activation via frizzled, dishevelled and axin functions. *Dev Camb Engl* 2008;135:367–75.
- [68] Cliffe A, Hamada F, Bienz M. A role of Dishevelled in relocating Axin to the plasma membrane during wingless signaling. *Curr Biol CB* 2003;13:960–6.
- [69] Cong F, Schweizer L, Varmus H. Wnt signals across the plasma membrane to activate the beta-catenin pathway by forming oligomers containing its receptors, Frizzled and LRP. *Development* 2004;131:5103–15.
- [70] Stamos JL, Weis WI. The β -catenin destruction complex. *Cold Spring Harb Perspect Biol* 2013;5.
- [71] Lee E, Salic A, Krüger R, Heinrich R, Kirschner MW. The roles of APC and Axin derived from experimental and theoretical analysis of the Wnt pathway. *PLoS Biol* 2003;1:E10.
- [72] Giles RH, van Es JH, Clevers H. Caught up in a Wnt storm: Wnt signaling in cancer. *Biochim Biophys Acta* 2003;1653:1–24.
- [73] Klaus A, Birchmeier W. Wnt signalling and its impact on development and cancer. *Nat Rev Cancer* 2008;8:387–98.
- [74] Chenn A, Walsh CA. Regulation of cerebral cortical size by control of cell cycle exit in neural precursors. *Science* 2002;297:365–9.
- [75] Chenn A, Walsh CA. Increased neuronal production, enlarged forebrains and cytoarchitectural distortions in beta-catenin overexpressing transgenic mice. *Cereb Cortex* 2003;13:599–606.
- [76] Cox RT, Kirkpatrick C, Peifer M. Armadillo is required for adherens junction assembly, cell polarity, and morphogenesis during drosophila embryogenesis. *J Cell Biol* 1996;134:133–48.
- [77] Peifer M, Polakis P. Wnt signaling in oncogenesis and embryogenesis—a look outside the nucleus. *Science* 2000;287:1606–9.
- [78] Song JL, Nigam P, Tektas SS, Selva E. microRNA regulation of Wnt signaling pathways in development and disease. *Cell Signal* 2015;27:1380–91.
- [79] Triboulet R, Mari B, Lin Y-L, Chable-Bessia C, Bennasser Y, Lebrigand K, et al. Suppression of MicroRNA-silencing pathway by HIV-1 during virus replication. *Science* 2007;315:1579–82.
- [80] Wang F-Z, Weber F, Croce C, Liu C-G, Liao X, Pellett PE. Human cytomegalovirus infection alters the expression of cellular MicroRNA species that affect its replication. *J Virol* 2008;82:9065–74.
- [81] Peng X, Li Y, Walters K-A, Rosenzweig ER, Lederer SL, Aicher LD, et al. Computational identification of hepatitis C virus associated microRNA-mRNA regulatory modules in human livers. *BMC Genomics* 2009;10:373.
- [82] Smith JL, Grey FE, Uhrlaub JL, Nikolich-Zugich J, Hirsch AJ. Induction of the cellular MicroRNA, Hs_154, by West Nile virus contributes to virus-mediated apoptosis through repression of antiapoptotic factors. *J Virol* 2012;86:5278–87.
- [83] Slonchak A, Shannon RP, Pali G, Khromykh AA. Human microRNA miR-532-5p exhibits antiviral activity against west Nile virus via suppression of host genes SESTD1 and TAB3 required for virus replication. *J Virol* 2016;90:2388–402.
- [84] Smith JL, Jeng S, McWeeny SK, Hirsch AJ. A microRNA screen identifies the WNT signaling pathway as a regulator of the interferon response during flavivirus infection. *J Virol* 2017;91.
- [85] Yang P, An H, Liu X, Wen M, Zheng Y, Rui Y, et al. The cytosolic nucleic acid sensor LRRFIP1 mediates the production of type I interferon via a β -catenin-dependent pathway. *Nat Immunol* 2010;11:487–94.
- [86] Zhu J, Coyne CB, Sarkar SN. PKC α regulates Sendai virus-mediated interferon induction through HDAC6 and β -catenin. *EMBO J* 2011;30:4838–49.
- [87] Baril M, Es-Saad S, Chatel-Chaix L, Fink K, Pham T, Raymond V-A, et al. Genome-wide RNAi screen reveals a new role of a WNT/CTNBB1 signaling pathway as negative regulator of virus-induced innate immune responses. *PLOS Pathog* 2013;9:e1003416.
- [88] Hillesheim A, Nordhoff C, Boergeling Y, Ludwig S, Wixler V. β -catenin promotes the type I IFN synthesis and the IFN-dependent signaling response but is suppressed by influenza A virus-induced RIG-I/NF- κ B signaling. *Cell Commun Signal* 2014;12:29.
- [89] Qin Y, Liu Q, Tian S, Xie W, Cui J, Wang R-F. TRIM9 short isoform preferentially promotes DNA and RNA virus-induced production of type I interferon by recruiting GSK3 β to TBK1. *Cell Res* 2016;26:613–28.
- [90] McLean E, Bhattarai R, Hughes BW, Mahalingam K, Bagasra O. Computational identification of mutually homologous Zika virus miRNAs that target microcephaly genes. *Libyan J Med* 2017;12.
- [91] Lopez-Valenzuela M, Ramírez O, Rosas A, García-Vargas S, de la Rasilla M, Lalueza-Fox C, et al. An Ancestral miR-1304 allele present in Neanderthals regulates genes involved in enamel formation and could explain dental differences with modern humans. *Mol Biol Evol* 2012;29:1797–806.
- [92] Trimborn M, Bell SM, Felix C, Rashid Y, Jafri H, Griffiths PD, et al. Mutations in microcephalin cause aberrant regulation of chromosome condensation. *Am J Hum Genet* 2004;75:261–6.
- [93] Xu X, Lee J, Stern DF. Microcephalin is a DNA damage response protein involved in regulation of CHK1 and BRCA1. *J Biol Chem* 2004;279:34091–4.
- [94] Trimborn M, Ghani M, Walther DJ, Dopatka M, Dutranoy V, Busche A, et al. Establishment of a mouse model with misregulated chromosome condensation due to defective Mcph1 function. *PLoS One* 2010;5:e9242.
- [95] Bilgüvar K, Öztürk AK, Louvi A, Kwan KY, Choi M, Tatli B, et al. Whole-exome sequencing identifies recessive WDR62 mutations in severe brain malformations. *Nature* 2010;467:207–10.
- [96] Yu TW, Mochida GH, Tischfield DJ, Sgaier SK, Flores-Sarnat L, Sergi CM, et al. Mutations in WDR62, encoding a centrosome-associated protein, cause microcephaly with simplified gyri and abnormal cortical architecture. *Nat Genet* 2010;42:1015–20.
- [97] Lucas EP, Raff JW. Maintaining the proper connection between the centrioles and the pericentriolar matrix requires Drosophila centrosomin. *J Cell Biol* 2007;178:725–32.
- [98] Graser S, Stierhof Y-D, Nigg EA. Cep68 and Cep215 (Cdk5rap2) are required for centrosome cohesion. *J Cell Sci* 2007;120:4321–31.
- [99] Zhang X, Liu D, Lv S, Wang H, Zhong X, Liu B, et al. CDK5RAP2 is required for spindle checkpoint function. *Cell Cycle* 2009;8:1206–16.
- [100] Kiyomitsu T, Obuse C, Yanagida M. Human Blinkin/AF15q14 is required for chromosome alignment and the mitotic checkpoint through direct interaction with Bub1 and BubR1. *Dev Cell* 2007;13:663–76.
- [101] Cheeseman IM, Hori T, Fukagawa T, Desai A. KNL1 and the CENP-H/I/K complex coordinately direct kinetochore assembly in vertebrates. *Mol Biol Cell* 2008;19:587–94.
- [102] Guernsey DL, Jiang H, Hussin J, Arnold M, Bouyakdan K, Perry S, et al. Mutations in centrosomal protein CEP152 in primary microcephaly families linked to MCPH4. *Am J Hum Genet* 2010;87:40–51.
- [103] do Carmo Avides M, Glover DM. Abnormal spindle protein, Asp, and the integrity of mitotic centrosomal microtubule organizing centers. *Science* 1999;283:1733–5.
- [104] Riparbelli MG, Callaini G, Glover DM, Avides Mdo C. A requirement for the abnormal spindle protein to organise microtubules of the central spindle for cytokinesis in Drosophila. *J Cell Sci* 2002;115:913–22.
- [105] Kouprina N, Pavlicek A, Collins NK, Nakano M, Noskov VN, Ohzeki J-I, et al. The microcephaly ASPM gene is expressed in proliferating tissues and encodes for a mitotic spindle protein. *Hum Mol Genet* 2005;14:2155–65.
- [106] Kirkham M, Müller-Reichert T, Oegema K, Grill S, Hyman AA. SAS-4 is a C. elegans centriolar protein that controls centrosome size. *Cell* 2003;112:575–87.
- [107] Hung L-Y, Chen H-L, Chang C-W, Li B-R, Tang TK. Identification of a novel microtubule-destabilizing motif in CPAP that binds to tubulin heterodimers and inhibits microtubule assembly. *Mol Biol Cell* 2004;15:2697–706.
- [108] Cho J-H, Chang C-J, Chen C-Y, Tang TK. Depletion of CPAP by RNAi disrupts centrosome integrity and induces multipolar spindles. *Biochem Biophys Res Commun* 2006;339:742–7.
- [109] Campaner S, Kaldis P, Izraeli S, Kirsch IR. Sil phosphorylation in a Pin1 binding domain affects the duration of the spindle checkpoint. *Mol Cell Biol* 2005;25:6660–72.
- [110] Pfaff KL, Straub CT, Chiang K, Bear DM, Zhou Y, Zon LI. The zebra fish cassiopeia mutant reveals that SIL is required for mitotic spindle organization. *Mol Cell Biol* 2007;27:5887–97.
- [111] Kumar A, Girimaji SC, Duvvari MR, Blanton SH. Mutations in STIL, encoding a pericentriolar and centrosomal protein, cause primary microcephaly. *Am J Hum Genet* 2009;84:286–90.
- [112] Blachon S, Gopalakrishnan J, Omori Y, Polyanovsky A, Church A, Nicastro D, et al. Drosophila asterless and vertebrate Cep152 Are orthologs essential for centriole duplication. *Genetics* 2008;180:2081–94.
- [113] Mirzaa GM, Vitre B, Carpenter G, Abramowicz I, Gleeson JG, Paciorkowski AR, et al. Mutations in CENPE define a novel kinetochore-centromeric mechanism for microcephalic primordial dwarfism. *Hum Genet* 2014;133:1023–39.
- [114] Strnad P, Leidel S, Vinogradova T, Euteneuer U, Khodjakov A, Gönczy P. Regulated HsSAS-6 levels ensure formation of a single procentriole per centriole during the centrosome duplication cycle. *Dev Cell* 2007;13:203–13.
- [115] Tang C-JC, Lin S-Y, Hsu W-B, Lin Y-N, Wu C-T, Lin Y-C, et al. The human microcephaly protein STIL interacts with CPAP and is required for procentriole formation. *EMBO J* 2011;30:4790–804.

- [116] Ben-Zvi A, Lacoste B, Kur E, Andreone BJ, Mayshar Y, Yan H, et al. Mfsd2a is critical for the formation and function of the blood-brain barrier. *Nature* 2014;509:507–11.
- [117] Alakbarzade V, Hameed A, Quek DQY, Chioza BA, Baple EL, Cazenave-Gassiot A, et al. A partially inactivating mutation in the sodium-dependent lysophosphatidylcholine transporter MFSD2A causes a non-lethal microcephaly syndrome. *Nat Genet* 2015;47:814–7.
- [118] Yamamoto S, Jaiswal M, Charng W-L, Gambin T, Karaca E, Mirzaa G, et al. A drosophila genetic resource of mutants to study mechanisms underlying human genetic diseases. *Cell* 2014;159:200–14.
- [119] Kaufmann T, Kukolj E, Brachner A, Beltzung E, Bruno M, Kostrhon S, et al. SIRT2 regulates nuclear envelope reassembly through ANKLE2 deacetylation. *J Cell Sci* 2016;129:4607–21.
- [120] Brown NJ, Marjanović M, Lüders J, Stracker TH, Costanzo V. Cep63 and cep152 cooperate to ensure centriole duplication. *PLoS One* 2013;8, e69986.
- [121] Alsara M, Löffler H, Fechter A, Bartek J, Krämer A. Cep63 recruits cdk1 to the centrosome-letter. *Cancer Res* 2015;75:777–8.
- [122] Najas S, Arranz J, Lochhead PA, Ashford AL, Oxley D, Delabar JM, et al. DYRK1A-mediated cyclin D1 degradation in neural stem cells contributes to the neurogenic cortical defects in down syndrome. *EBioMedicine* 2015;2:120–34.
- [123] Ji J, Lee H, Argiropoulos B, Dorrani N, Mann J, Martinez-Agosto JA, et al. DYRK1A haploinsufficiency causes a new recognizable syndrome with microcephaly, intellectual disability, speech impairment, and distinct facies. *Eur J Hum Genet* 2015;23:1473–81.
- [124] Russ AP, Wattler S, Colledge WH, Aparicio SA, Carlton MB, Pearce JJ, et al. Eomesodermin is required for mouse trophoblast development and mesoderm formation. *Nature* 2000;404:95–9.
- [125] Arnold SJ, Huang G-J, Cheung AFP, Era T, Nishikawa S-I, Bikoff EK, et al. The T-box transcription factor Eomes/Tbr2 regulates neurogenesis in the cortical subventricular zone. *Genes Dev* 2008;22:2479–84.
- [126] Sessa A, Mao C, Hadjantonakis A-K, Klein WH, Broccoli V. Tbr2 directs conversion of radial glia into basal precursors and guides neuronal amplification by indirect neurogenesis in the developing neocortex. *Neuron* 2008;60:56–69.
- [127] Rapley J, Nicolàs M, Groen A, Regué L, Bertran MT, Caelles C, et al. The NIMA-family kinase Nek6 phosphorylates the kinesin Eg5 at a novel site necessary for mitotic spindle formation. *J Cell Sci* 2008;121:3912–21.
- [128] Fahnrikar A, Tole S, Baas PW. Kinesin-5, a mitotic microtubule-associated motor protein, modulates neuronal migration. *Mol Biol Cell* 2011;22:1561–74.
- [129] Yu X, Chen J. DNA damage-induced cell cycle checkpoint control requires CtIP, a phosphorylation-dependent binding partner of BRCA1 C-terminal domains. *Mol Cell Biol* 2004;24:9478–86.
- [130] You Z, Shi LZ, Zhu Q, Wu P, Zhang Y-W, Basilio A, et al. CtIP Links DNA double-strand break sensing to resection. *Mol Cell* 2009;36:954–69.
- [131] Manning AL, Dyson NJ. RB: mitotic implications of a tumour suppressor. *Nat Rev Cancer* 2012;12:220–6.
- [132] Clausen TH, Lamark T, Isakson P, Finley K, Larsen KB, Brech A, et al. p62/SQSTM1 and ALFY interact to facilitate the formation of p62 bodies/ALIS and their degradation by autophagy. *Autophagy* 2010;6:330–44.
- [133] Wu DJ, Gu R, Sarin R, Zavodovskaya R, Chen C-P, Christiansen BA, et al. Autophagy-linked FYVE containing protein WDFY3 interacts with TRAF6 and modulates RANKL-induced osteoclastogenesis. *J Autoimmun* 2016;73:73–84.