



Case Report

Prenatal diagnosis and molecular cytogenetic characterization of an interstitial deletion of 18q12.1-q12.3 encompassing *DTNA*, *CELF4* and *SETBP1*

Chih-Ping Chen^{a, b, c, d, e, f, g, *}, Chih-Heng Hsieh^h, Schu-Rern Chern^b, Peih-Shan Wuⁱ, Shin-Wen Chen^a, Shih-Ting Lai^a, Tzu-Yun Chuang^a, Chien-Wen Yang^b, Chen-Chi Lee^a, Wayseen Wang^{b, j}

^a Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^b Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^c Department of Medicine, Mackay Medical College, New Taipei, Taiwan

^d Department of Biotechnology, Asia University, Taichung, Taiwan

^e School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^f Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^g Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^h Department of Obstetrics and Gynecology, BIN KUN Women's & Children's Hospital, Taoyuan, Taiwan

ⁱ Gene Biodesign Co. Ltd, Taipei, Taiwan

^j Department of Bioengineering, Tatung University, Taipei, Taiwan

ARTICLE INFO

Article history:

Accepted 17 October 2017

Keywords:

18q12.1-q12.3 deletion

CELF4

DTNA

Prenatal diagnosis

SETBP1

ABSTRACT

Objective: We present prenatal diagnosis and molecular cytogenetic characterization of an interstitial deletion of 18q12.1-q12.3.

Case report: A 35-year-old woman underwent amniocentesis at 18 weeks of gestation because of advanced maternal age. Amniocentesis revealed a karyotype of 46,XX,del(18)(q12.1q12.3). The fetal ultrasound was unremarkable. The woman underwent repeat amniocentesis at 20 weeks of gestation. Array comparative genomic hybridization (aCGH) using uncultured amniocytes revealed a 10.76-Mb interstitial deletion 18q12.1-q12.3 or arr 18q12.1q12.3 (31,944,347–42,704,784) × 1.0 encompassing 19 Online Mendelian Inheritance of Man (OMIM) genes including *DTNA*, *CELF4* and *SETBP1*. Metaphase fluorescence *in situ* hybridization analysis on cultured amniocytes confirmed an 18q proximal interstitial deletion. The parental karyotypes were normal. Polymorphic DNA marker analysis determined a paternal origin of the deletion. The pregnancy was subsequently terminated at 24 weeks of gestation, and a 650-g fetus was delivered with characteristic facial dysmorphism.

Conclusion: aCGH analysis and polymorphic DNA marker analysis at amniocentesis are useful for determination of the deleted genes and the parental origin of the *de novo* deletion, and the acquired information is helpful for genetic counseling.

© 2017 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Amniocentesis because of advanced maternal age or abnormal ultrasound may incidentally detect a paternal origin *de novo*

chromosome aberration [1–3]. We previously reported prenatal diagnosis of a case with 18q12.1-q12.3 interstitial deletion [1]. Here, we present an additional case with an 18q12.1-q12.3 interstitial deletion.

Case report

A 35-year-old, gravida 2, para 1, woman underwent amniocentesis at 18 weeks of gestation because of advanced maternal age. Her husband was 31 years old, and there was no family history of

* Corresponding author. Department of Obstetrics and Gynecology, MacKay Memorial Hospital 92, Section 2, Chung-Shan North Road, Taipei, 10449, Taiwan. Fax: +886 2 25433642, +886 2 25232448.

E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).

congenital malformations. During this pregnancy, amniocentesis revealed a karyotype of 46,XX,del(18)(q12.1q12.3). The fetal ultrasound was unremarkable. The parental karyotypes were normal. The parents consulted our department and requested for repeat amniocentesis. Repeat amniocentesis was performed at 20 weeks of gestation. aCGH analysis on uncultured amniocytes by SurePrint G3 Unrestricted CGH ISCA v2, 8 × 60 K Array (Agilent Technologies, Santa Clara, CA, USA) revealed a 10.76-Mb interstitial deletion of 18q12.1–q12.3 or arr 18q12.1q12.3 (31,944,347–42,704,784) × 1.0 [GRCh37 (hg19)] encompassing 19 Online Mendelian Inheritance of in Man (OMIM) genes including *DTNA*, *CELF4* and *SETBP1* (Fig. 1). Conventional cytogenetic analysis on cultured amniocytes revealed a result of 46,XX,del(18)(q12.1q12.3) (Fig. 2). Polymorphic DNA marker analysis using the DNAs extracted from parental bloods and uncultured amniocytes determined a paternal origin of the deletion (Fig. 3). Metaphase fluorescence *in situ* hybridization analysis on cultured amniocytes confirmed an 18q proximal interstitial deletion (Fig. 4). The pregnancy was subsequently terminated at 24

weeks of gestation, and a 650-g fetus was delivered with characteristic facial features of telecanthus, frontal bossing, deep-set eyes and midface hypoplasia (Fig. 5).

Discussion

Proximal interstitial 18q deletions encompassing the bands 18q12→q21 define a rare del(18)(q12.1q12.3) syndrome with characteristic facial features of telecanthus, frontal bossing, deep-set eyes and midface hypoplasia, behavior problems, autism spectrum disorder, hyperactivity, aggressivity, moderate to severe mental retardation, hypotonia, seizures and language impairment but lack of major congenital defects [4–14]. The present case had a 10.76-Mb interstitial deletion 18q12.1–q12.3 encompassing *DTNA*, *CELF4* and *SETBP1*.

SETBP1 (OMIM 611060) encodes set-binding protein 1 and is located at 18q12.3. Mutations or deletions of *SETBP1* can be associated with autosomal dominant mental retardation-29 (OMIM




Fig. 1. Array comparative genomic hybridization on uncultured amniocytes shows a 10.76-Mb interstitial deletion of 18q12.1–q12.3 encompassing *DTNA*, *CELF4* and *SETBP1*. (A) and (B) Chromosome 18 zoom-in views.




Fig. 2. A karyotype 46,XX,del(18)(q12.1q12.3).




Fig. 3. Polymorphic DNA marker analysis using the DNAs extracted from parental bloods and uncultured amniocytes with the informative markers of D18S976 (18p11.31), D18S865 (18q12.3) and D18S868 (18q22.1) shows that the fetus has biparental inheritance in D18S976 and D18S868. However, in D18S865 at 18q12.3, the fetus inherits only one maternal allele, indicating a paternal origin of the deletion.

616078), and heterozygous *de novo* mutations in *SETBP1* cause autosomal dominant Schinzel-Giedion midface retraction syndrome (OMIM 269150). Bouquillon et al. [15] reported two patients with an 18q12.3 deletion, *SETBP1* haploinsufficiency and

expressive speech delay. Filges et al. [16] reported *SETBP1* haploinsufficiency in two patients with an 18q12.3 microdeletion, global developmental and expressive language delay and minor facial anomalies. Marseglia et al. [17] reported a patient with an




Fig. 4. Metaphase fluorescence *in situ* hybridization analysis on cultured amniocytes using the 18q12.2-specific probe of RP11-938M23 [fluorescein isothiocyanate (FITC), spectrum green] and the 18p11.22-specific probe of RP11-918F20 (Texas red, spectrum red) shows both the green and red signals in a normal chromosome 18, and only the red signal in the chromosome of del(18), indicating an 18q proximal interstitial deletion in the del(18). del = deletion.

18q12.3 microdeletion causing *SETBP1* haploinsufficiency in association with mild mental retardation and expressive speech impairment. Rauch et al. [18] and Coe et al. [19] reported patients with loss-of-function frameshift or nonsense mutations in *SETBP1* causing speech and motor delays, mild dysmorphic features and behavioral difficulties. Heterozygous *de novo* mutations in *SETBP1* cause Schinzel-Giedion midface retraction syndrome which is characterized by severe mental retardation, distinctive facial features of midface retraction, skeletal abnormalities, genitourinary

and renal malformations, cardiac defects and neuroepithelial neoplasia [20–28].

CEL4 (OMIM 612679) located at 18q12.2 is highly expressed in brain and gallbladder [29]. Halgren et al. [30] reported haploinsufficiency of *CEL4* at 18q12.2 in a patient with developmental and behavioral disorders, myopia, obesity and febrile seizures in childhood, and suggested that *CEL4* is important for human brain development. Barone et al. [31] reported familial 18q12.2 microdeletion involving *CEL4* in a child with syndromic intellectual disability and autism and her mother with minor dysmorphism, mild intellectual disability and autistic behavior, and suggested that *CEL4* plays a role in brain development and autism spectrum disorders. 18q12.2 deletion syndrome is a highly recognized condition characterized by the neuropsychiatric phenotype of developmental delay, intellectual disability, seizures, motor coordination disorder, behavioral disturbances and autism spectrum disorders, and other features of facial dysmorphism, myopia and obesity [8,13,30,32].

DTNA (OMIM 601239) located at 18q12.1 and encodes α -dystrobrevin which participates in the formation of sarcoglycan–sarcospan complex [33]. Wang et al. [34] identified a 4-Mb *de novo* duplication of 18q12.1 in a girl with autistic disorder, intellectual disability, focal epilepsy and idiopathic small stature and suggested that deletion or duplication of *DTNA* at 18q12.1 is associated with physical and neurological development including autism and intellectual disability.

Poissonnier et al. [35] reported two unrelated mentally retarded patients with del(18)(q12.1q12.3) that manifested facial dysmorphism, small midface, prominent forehead, epicanthic folds, full periorbital tissue, wide/flat nasal bridge, abnormal ears, behavioral disorders, autistic-like disorders, hypotonia and seizures. Our case is an additional case of del(18)(q12.1q12.3). Our case shows that array comparative genomic hybridization (aCGH) analysis and polymorphic DNA marker analysis at amniocentesis are useful for determination of the deleted genes and the parental origin of the *de novo* deletion, and the acquired information is helpful for genetic counseling.



Fig. 5. Anterior and lateral views of the craniofacial appearance of the proband at birth.

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

Acknowledgements

This work was supported by research grants MOST-105-2314-B-195-012 from the Ministry of Science and Technology and MMH-E-106-04 from MacKay Memorial Hospital, Taipei, Taiwan.

References

- [1] Chen C-P, Huang M-C, Chen Y-Y, Chern S-R, Wu P-S, Chen Y-T, et al. Prenatal diagnosis of de novo interstitial deletions involving 5q23.1-q23.3 and 18q12.1-q12.3 by array CGH using uncultured amniocytes in a pregnancy with fetal interrupted aortic arch and atrial septal defect. *Gene* 2013;531:496–501.
- [2] Chen C-P, Chen C-Y, Chern S-R, Wu P-S, Chen Y-N, Chen S-W, et al. Prenatal diagnosis and molecular cytogenetic characterization of a de novo 4.858-Mb microdeletion in 15q14 associated with *ACTC1* and *MEIS2* haploinsufficiency and tetralogy of Fallot. *Taiwan J Obstet Gynecol* 2016;55:270–4.
- [3] Chen C-P, Hung F-Y, Chern S-R, Wu P-S, Chen Y-N, Chen S-W, et al. Prenatal diagnosis and molecular cytogenetic characterization of de novo partial monosomy 3p (3p26.3→pter) and partial trisomy 16q (16q23.1→qter). *Taiwan J Obstet Gynecol* 2016;55:288–92.
- [4] Chudley AE, Bauder F, Ray M, McAlpine PJ, Pena SDJ, Hamerton JL. Familial mental retardation in a family with an inherited chromosome rearrangement. *J Med Genet* 1974;11:353–66.
- [5] Wilson GN, Al Saadi AA. Obesity and abnormal behavior associated with interstitial deletion of chromosome 18 (q12.2q21.1). *J Med Genet* 1989;26:62–3.
- [6] Schinzel A, Binkert F, Lillington DM, Sands M, Stocks RJ, Lindenbaum RH, et al. Interstitial deletion of the long arm of chromosome 18, del(18)(q12.2q21.1): a report of three cases of an autosomal deletion with a mild phenotype. *J Med Genet* 1991;28:352–5.
- [7] Engelen JJM, Loots WJG, Albrechts JCM, Plomp AS, Van Der Meer SB, Vies JSH, et al. Characterization of a de novo unbalanced translocation t(14q18q) using microdissection and fluorescence *in situ* hybridization. *Am J Med Genet* 1998;75:409–13.
- [8] McEntagart M, Carey A, Breen C, McQuaid S, Stallings RL, Green AJ, et al. Molecular characterisation of a proximal chromosome 18q deletion. *J Med Genet* 2001;38:128–9.
- [9] Tinkle BT, Christianson CA, Schorry EK, Webb T, Hopkin RJ. Long-term survival in a patient with del(18)(q12.2q21.1). *Am J Med Genet* 2003;119A:66–70.
- [10] Gribble SM, Prigmore E, Burford DC, Porter KM, Ng BL, Douglas EJ, et al. The complex nature of constitutional de novo apparently balanced translocations in patients presenting with abnormal phenotypes. *J Med Genet* 2005;42:8–16.
- [11] Kotzot D, Haberlandt E, Fauth C, Baumgartner S, Scholl-Bürgi S, Utermann G. Del(18)(q12.2q21.1) caused by a paternal sister chromatid rearrangement in a developmentally delayed girl. *Am J Med Genet* 2005;135A:304–7.
- [12] Cody JD, Sebold C, Malik A, Heard P, Carter E, Crandall A, et al. Recurrent interstitial deletions of proximal 18q: a new syndrome involving expressive speech delay. *Am J Med Genet* 2007;143A:1181–90.
- [13] Feenstra I, Vissers LELM, Orsel M, van Kessel AG, Brunner HG, Veltman JA, et al. Genotype–phenotype mapping of chromosome 18q deletions by high-resolution array CGH: an update of the phenotypic map. *Am J Med Genet* 2007;143A:1858–67.
- [14] Buysse K, Menten B, Oostra A, Tavernier S, Mortier GR, Speleman F. Delineation of a critical region on chromosome 18 for the del(18)(q12.2q21.1) syndrome. *Am J Med Genet* 2008;146A:1330–4.
- [15] Bouquillon S, Andrieux J, Landais E, Duban-Bedu B, Boidein F, Lenne B, et al. A 5.3Mb deletion in chromosome 18q12.3 as the smallest region of overlap in two patients with expressive speech delay. *Eur J Med Genet* 2011;54:194–7.
- [16] Filges I, Shimojima K, Okamoto N, Rothlisberger B, Weber P, Huber AR, et al. Reduced expression by *SETBP1* haploinsufficiency causes developmental and expressive language delay indicating a phenotype distinct from Schinzel-Giedion syndrome. *J Med Genet* 2011;48:117–22.
- [17] Marseglia G, Scordo MR, Pescucci C, Nannetti G, Biagini E, Scandurra V, et al. 372 kb microdeletion in 18q12.3 causing *SETBP1* haploinsufficiency associated with mild mental retardation and expressive speech impairment. *Eur J Med Genet* 2012;55:216–21.
- [18] Rauch A, Wiczorek D, Graf E, Wieland T, Ende S, Schwarzmayr T, et al. Range of genetic mutations associated with severe non-syndromic sporadic intellectual disability: an exome sequencing study. *Lancet* 2012;380:1674–82.
- [19] Coe BP, Witherspoon K, Rosenfeld JA, van Bon BWM, Vulto-van Silfhout AT, Bosco P, et al. Refining analyses of copy number variation identifies specific genes associated with developmental delay. *Nat Genet* 2014;46:1063–71.
- [20] Schinzel A, Giedion A. A syndrome of severe midface retraction, multiple skull anomalies, clubfeet, and cardiac and renal malformations in sibs. *Am J Med Genet* 1978;1:361–75.
- [21] Schinzel A. A syndrome of midface retraction, multiple radiological anomalies, renal malformations and hypertrichosis. *Hum Genet* 1982;62:382 (Letter).
- [22] Al-Gazali LI, Farndon PA, Burn J, Massey R, Mueller RF. The Schinzel-Giedion syndrome: clinical features and natural history. *J Med Genet* 1988;25:275 [Abstract].
- [23] Al-Gazali LI, Farndon P, Burn J, Flannery DB, Davison C, Mueller RF. The Schinzel-Giedion syndrome. *J Med Genet* 1990;27:42–7.
- [24] Hoischen A, van Bon BWM, Gilissen C, Arts P, van Lier B, Steehouwer M, et al. De novo mutations of *SETBP1* cause Schinzel-Giedion syndrome. *Nat Genet* 2010;42:483–5.
- [25] Suphapeetiporn K, Srichomthong C, Shotelersuk V. *SETBP1* mutations in two Thai patients with Schinzel-Giedion syndrome. *Clin Genet* 2011;79:391–3.
- [26] Ko JM, Lim BC, Kim KJ, Hwang YS, Ryu HW, Lee JH, et al. Distinct neurological features in a patient with Schinzel-Giedion syndrome caused by a recurrent *SETBP1* mutation. *Childs Nerv Syst* 2013;29:525–9.
- [27] Carvalho E, Honjo R, Magalhães M, Yamamoto G, Rocha K, Naslavsky M, et al. Schinzel-Giedion syndrome in two Brazilian patients: report of a novel mutation in *SETBP1* and literature review of the clinical features. *Am J Med Genet* 2015;167A:1039–46.
- [28] Takeuchi A, Okamoto N, Fujinaga S, Morita H, Shimizu J, Akiyama T, et al. Progressive brain atrophy in Schinzel-Giedion syndrome with a *SETBP1* mutation. *Eur J Med Genet* 2015;58:369–71.
- [29] Ladd AN, Nguyen NH, Malhotra K, Cooper TA. CELF6, a member of the CELF family of RNA-binding proteins, regulates muscle-specific splicing enhancer-dependent alternative splicing. *J Biol Chem* 2004;279:17756–64.
- [30] Halgren C, Bache I, Bak M, Myatt MW, Anderson CM, Brøndum-Nielsen K, et al. Haploinsufficiency of *CELF4* at 18q12.2 is associated with developmental and behavioral disorders, seizures, eye manifestations, and obesity. *Eur J Hum Genet* 2012;20:1315–9.
- [31] Barone R, Fichera M, De Grandi M, Battaglia M, Lo Faro V, Mattina T, et al. Familial 18q12.2 deletion supports the role of RNA-binding protein *CELF4* in autism spectrum disorders. *Am J Med Genet* 2017;173A:1649–55.
- [32] Gilling M, Lauritsen MB, Møller M, Henriksen KF, Vicente A, Oliveira G, et al. A 3.2 Mb deletion on 18q12 in a patient with childhood autism and high-grade myopia. *Eur J Hum Genet* 2008;16:312–9.
- [33] Yoshida M, Hama H, Ishikawa-Sakurai M, Imamura M, Mizuno Y, Araiishi K, et al. Biochemical evidence for association of dystrobrevin with the sarcoglycan-sarcospan complex as a basis for understanding sarcoglycanopathy. *Hum Mol Genet* 2000;9:1033–40.
- [34] Wang P, Carrion P, Qiao Y, Tyson C, Hrynychak M, Calli K, et al. Genotype-phenotype analysis of 18q12.1-q12.2 copy number variation in autism. *Eur J Med Genet* 2013;56:420–5.
- [35] Poissonnier M, Turleau C, Olivier-Martin M, Milleret-Proyart MJ, Prieur M, Dubos M, et al. Interstitial deletion of the proximal region of the long arm of chromosome 18, del(18q12) a distinct clinical entity? A report of two new cases. *Ann Genet* 1992;35:146–51.